

Protecting yourself against COVID-19

COVID-19 VACCINATION SCHEDULE

Stay up to date on your vaccinations with a primary series and a booster.

Pfizer-BioNTech

5-11 years (pediatric formulation)



Pfizer-BioNTech

12 and older (adult/adolescent formulation)




Moderna

18 and older



Johnson & Johnson (Janssen)**

18 and older



You should get a second booster if you are at higher risk for serious illness from COVID-19.

People in certain groups or situations are now eligible to get a 2nd COVID-19 booster four months after the 1st booster.

An extra booster can strengthen protection for people who are at higher risk for serious illness, hospitalization, or death. Your healthcare provider can help you review your options.

*Recommended intervals between doses can vary based upon age and other risk factors. Talk to your healthcare provider about the best timing between the first and second dose for your situation. An 8-week interval may be optimal for people ages 12-64 years, and especially for males ages 12-39 years. A shorter interval (3 weeks for Pfizer-BioNTech; 4 weeks for Moderna) between the first and second dose remains the recommended interval for adults ages 65 years and older and others who need early protection due to increased risk.

**The COVID-19 mRNA vaccines (Pfizer or Moderna) should be given in most cases. The Johnson & Johnson (Janssen) vaccine is authorized for limited uses. People age 18 and older may receive the Johnson & Johnson vaccine if they are unable to receive an mRNA vaccine due to allergies or severe reactions to the vaccine or its ingredients OR if they would otherwise not receive a COVID-19 vaccine.

*** 2nd boosters can only be Pfizer or Moderna. People ages 12-17 can only receive Pfizer.

