

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 10

Counties: Allen, Auglaize, Champaign, Hancock, Hardin, Logan, Mercer, Paulding, Putnam, Shelby, Van Wert

As of June 30, 2019

Reported new diagnoses of HIV infection in 2018 by disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	2018 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	5.1	13	68%	11	73%	1	33%	1	100%
Females	2.3	6	32%	4	27%	2	67%	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	*	2	11%	1	7%	-	-	1	100%
25-29	15.6	5	26%	4	27%	1	33%	-	-
30-34	*	4	21%	3	20%	1	33%	-	-
35-39	*	-	-	-	-	-	-	-	-
40-44	*	3	16%	3	20%	-	-	-	-
45-49	*	1	5%	1	7%	-	-	-	-
50-54	*	-	-	-	-	-	-	-	-
55-64	*	3	16%	2	13%	1	33%	-	-
65+	*	1	5%	1	7%	-	-	-	-
Race/Ethnicity ^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	33.4	6	32%	4	27%	1	33%	1	100%
Hispanic/Latinx	*	-	-	-	-	-	-	-	-
White	2.8	13	68%	11	73%	2	67%	-	-
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity ^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	*	4	21%	3	20%	-	-	1	100%
Black/African-American Females	*	2	11%	1	7%	1	33%	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	3.9	9	47%	8	53%	1	33%	-	-
White Females	*	4	21%	3	20%	1	33%	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	3.7	19		15		3		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	6	46%	5	45%	-	-	1	100%
Injection drug use (IDU)	1	8%	1	9%	-	-	-	-
Male-to-male sexual contact and IDU	1	8%	1	9%	-	-	-	-
Heterosexual contact	2	15%	1	9%	1	100%	-	-
Other/unknown	3	23%	3	27%	-	-	-	-
Subtotal	13	100%	11	100%	1	100%	1	100%
Female adult or adolescent								
Injection drug use	1	17%	1	25%	-	-	-	-
Heterosexual contact	4	67%	3	75%	1	50%	-	-
Other/unknown	1	17%	-	-	1	50%	-	-
Subtotal	6	100%	4	100%	2	100%	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	19		15		3		1	

Reported new diagnoses of HIV infection in 2018 by disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	32%	5	33%	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	6	32%	4	27%	2	67%	-	-
Male-to-male sexual contact & IDU	1	5%	1	7%	-	-	-	-
IDU & Heterosexual contact	2	11%	2	13%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	4	21%	3	20%	1	33%	-	-
Total	19		15		3		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 10

2018 diagnosis of HIV infection																		
Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx*			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	2	33%	*	-	-	*	-	-	*	-	-
25-29	*	-	-	*	-	-	*	1	17%	*	-	-	*	4	31%	*	-	-
30-34	*	-	-	*	-	-	*	1	17%	*	-	-	*	3	23%	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	3	23%	*	-	-
45-49	*	-	-	*	-	-	*	-	-	*	-	-	*	1	8%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	1	17%	*	-	-	*	2	15%	*	-	-
65+	*	-	-	*	-	-	*	1	17%	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	33.4	6		*	-	-	2.8	13		*	-	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

2018 diagnosis of HIV infection												
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	3	75%	-	-	3	33%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	11%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	1	11%	-	-
Heterosexual contact	-	-	-	-	1	25%	-	-	1	11%	-	-
Other/unknown	-	-	-	-	-	-	-	-	3	33%	-	-
Subtotal	-	-	-	-	4	100%	-	-	9	100%	-	-
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	-	-	1	25%	-	-
Heterosexual contact	-	-	-	-	1	50%	-	-	3	75%	-	-
Other/unknown	-	-	-	-	1	50%	-	-	-	-	-	-
Subtotal	-	-	-	-	2	100%	-	-	4	100%	-	-
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-		-		6		-		13		-	

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

Exposure Category ^d	2018 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	3	50%	-	-	3	23%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	2	33%	-	-	4	31%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	8%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	15%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	1	17%	-	-	3	23%	-	-
Total	-	-	-	-	6	-	-	-	13	-	-	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 10

Characteristic	Diagnosis of HIV Infection by Year															Cumulative Diagnoses	
	2014			2015			2016			2017			2018				
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	4.7	12	80%	9.0	23	79%	3.5	9	53%	4.7	12	71%	5.1	13	68%	493	78%
Females	*	3	20%	2.3	6	21%	3.1	8	47%	2.0	5	29%	2.3	6	32%	142	22%
Age at diagnosis (yr)																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	7	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
15-19	*	-	-	*	2	7%	*	-	-	*	1	6%	*	-	-	17	3%
20-24	*	3	20%	14.6	5	17%	*	1	6%	*	2	12%	*	2	11%	74	12%
25-29	*	4	27%	*	3	10%	16.5	5	29%	*	1	6%	15.6	5	26%	102	16%
30-34	*	2	13%	17.0	5	17%	*	3	18%	*	2	12%	*	4	21%	108	17%
35-39	*	2	13%	*	3	10%	*	2	12%	*	1	6%	*	-	-	90	14%
40-44	*	1	7%	*	4	14%	*	-	-	*	3	18%	*	3	16%	91	14%
45-49	*	1	7%	*	1	3%	*	1	6%	*	2	12%	*	1	5%	55	9%
50-54	*	-	-	*	3	10%	*	2	12%	14.6	5	29%	*	-	-	44	7%
55-64	*	2	13%	*	3	10%	*	3	18%	*	-	-	*	3	16%	36	6%
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	1	5%	9	1%
Race/Ethnicity^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Black/African-American	*	4	27%	*	2	7%	*	3	18%	*	-	-	33.4	6	32%	105	17%
Hispanic/Latinx	*	-	-	*	1	3%	*	1	6%	*	4	24%	*	-	-	24	4%
White	2.3	11	73%	5.4	25	86%	2.8	13	76%	2.6	12	71%	2.8	13	68%	485	76%
Multi-Race	*	-	-	*	1	3%	*	-	-	*	1	6%	*	-	-	20	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Race/Ethnicity^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	1	<1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American Males	*	3	20%	*	1	3%	*	1	6%	*	-	-	*	4	21%	62	10%
Black/African-American Females	*	1	7%	*	1	3%	*	2	12%	*	-	-	*	2	11%	43	7%
Hispanic/Latino Males	*	-	-	*	1	3%	*	1	6%	*	3	18%	*	-	-	15	2%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	1	6%	*	-	-	9	1%
White Males	3.9	9	60%	9.1	21	72%	3.0	7	41%	3.5	8	47%	3.9	9	47%	399	63%
White Females	*	2	13%	*	4	14%	2.6	6	35%	*	4	24%	*	4	21%	86	14%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	1	6%	*	-	-	16	3%
Multi-Race Females	*	-	-	*	1	3%	*	-	-	*	-	-	*	-	-	4	1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Total	2.9	15		5.6	29		3.3	17		3.3	17		3.7	19		635	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Diagnosis of HIV Infection by Year										Cumulative Diagnoses	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	50%	19	83%	5	56%	11	92%	6	46%	323	66%
Injection drug use (IDU)	1	8%	1	4%	1	11%	1	8%	1	8%	27	5%
Male-to-male sexual contact and IDU	1	8%	-	-	1	11%	-	-	1	8%	30	6%
Heterosexual contact	-	-	-	-	2	22%	-	-	2	15%	28	6%
Other/unknown	4	33%	3	13%	-	-	-	-	3	23%	85	17%
Subtotal	12	100%	23	100%	9	100%	12	100%	13	100%	493	100%
Female adult or adolescent												
Injection drug use	-	-	3	50%	4	50%	2	40%	1	17%	27	20%
Heterosexual contact	3	100%	3	50%	4	50%	2	40%	4	67%	99	73%
Other/unknown	-	-	-	-	-	-	1	20%	1	17%	10	7%
Subtotal	3	100%	6	100%	8	100%	5	100%	6	100%	136	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	5	83%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	17%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	15		29		17		17		19		635	

Trends in reported diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Diagnosis of HIV infection by Year										Cumulative Diagnoses	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	5	33%	18	62%	5	29%	10	59%	6	32%	297	47%
Injection drug use (IDU) only	1	7%	1	3%	-	-	2	12%	-	-	23	4%
Heterosexual contact only	3	20%	3	10%	6	35%	2	12%	6	32%	128	20%
Male-to-male sexual contact & IDU	1	7%	-	-	1	6%	-	-	1	5%	23	4%
IDU & Heterosexual contact	-	-	3	10%	5	29%	1	6%	2	11%	31	5%
Male-to-male sexual contact & Heterosexual contact	1	7%	1	3%	-	-	1	6%	-	-	26	4%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	4	27%	3	10%	-	-	1	6%	4	21%	95	15%
Total	15		29		17		17		19		635	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 10

Characteristic	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	5	63%	3	75%	2	50%
Females	3	38%	1	25%	2	50%
Age at AIDS diagnosis (yr)						
<13	-	-	-	-	-	-
13-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	1	13%	-	-	1	25%
25-29	1	13%	-	-	1	25%
30-34	2	25%	1	25%	1	25%
35-39	1	13%	1	25%	-	-
40-44	-	-	-	-	-	-
45-49	-	-	-	-	-	-
50-54	1	13%	1	25%	-	-
55-64	1	13%	-	-	1	25%
65+	1	13%	1	25%	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African-American	2	25%	-	-	2	50%
Hispanic/Latinx	-	-	-	-	-	-
White	6	75%	4	100%	2	50%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	1	13%	-	-	1	25%
Black/African-American Females	1	13%	-	-	1	25%
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	4	50%	3	75%	1	25%
White Females	2	25%	1	25%	1	25%
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	8		4		4	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 10

Transmission Category ^b	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3	60%	2	67%	1	50%
Injection drug use (IDU)	-	-	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	1	20%	-	-	1	50%
Other/unknown	1	20%	1	33%	-	-
Subtotal	5	100%	3	100%	2	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	2	67%	1	100%	1	50%
Other/unknown	1	33%	-	-	1	50%
Subtotal	3	100%	1	100%	2	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	8		4		4	

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 10

Exposure Category ^c	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	38%	2	50%	1	25%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	3	38%	1	25%	2	50%
Male-to-male sexual contact & IDU	-	-	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	2	25%	1	25%	1	25%
Total	8		4		4	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 10

Characteristic	Year of AIDS Diagnosis												Cumulative AIDS	
	2014		2015		2016		2017		2018					
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%		
Sex at birth														
Males	11	85%	8	67%	11	92%	10	91%	5	63%	339	83%		
Females	2	15%	4	33%	1	8%	1	9%	3	38%	71	17%		
Age at AIDS diagnosis (yr)														
<13	-	-	-	-	-	-	-	-	-	-	5	1%		
13-14	-	-	-	-	-	-	-	-	-	-	-	-		
15-19	-	-	-	-	-	-	-	-	-	-	4	1%		
20-24	1	8%	-	-	-	-	-	-	1	13%	14	3%		
25-29	1	8%	2	17%	2	17%	-	-	1	13%	61	15%		
30-34	3	23%	2	17%	1	8%	1	9%	2	25%	67	16%		
35-39	1	8%	2	17%	-	-	1	9%	1	13%	60	15%		
40-44	1	8%	-	-	-	-	2	18%	-	-	57	14%		
45-49	1	8%	3	25%	3	25%	2	18%	-	-	59	14%		
50-54	1	8%	1	8%	2	17%	4	36%	1	13%	40	10%		
55-64	3	23%	2	17%	4	33%	1	9%	1	13%	35	9%		
65+	1	8%	-	-	-	-	-	-	1	13%	8	2%		
Race/Ethnicity ^a														
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-		
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-		
Black/African-American	4	31%	4	33%	-	-	1	9%	2	25%	59	14%		
Hispanic/Latinx	-	-	-	-	1	8%	1	9%	-	-	11	3%		
White	8	62%	8	67%	11	92%	9	82%	6	75%	328	80%		
Multi-Race	1	8%	-	-	-	-	-	-	-	-	12	3%		
Race/Ethnicity ^a and Sex at birth														
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-		
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-		
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-		
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-		
Black/African-American Males	3	23%	1	8%	-	-	1	9%	1	13%	36	9%		
Black/African-American Females	1	8%	3	25%	-	-	-	-	1	13%	23	6%		
Hispanic/Latino Males	-	-	-	-	1	8%	1	9%	-	-	7	2%		
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	4	1%		
White Males	7	54%	7	58%	10	83%	8	73%	4	50%	286	70%		
White Females	1	8%	1	8%	1	8%	1	9%	2	25%	42	10%		
Multi-Race Males	1	8%	-	-	-	-	-	-	-	-	10	2%		
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	2	<1%		
Total	13		12		12		11		8		410			

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	4	36%	6	75%	7	64%	8	80%	3	60%	227	67%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	-	-	15	4%
Male-to-male sexual contact and IDU	1	9%	-	-	1	9%	-	-	-	-	17	5%
Heterosexual contact	1	9%	-	-	1	9%	1	10%	1	20%	20	6%
Other/unknown	5	45%	2	25%	2	18%	1	10%	1	20%	60	18%
Subtotal	11	100%	8	100%	11	100%	10	100%	5	100%	339	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	100%	1	100%	-	-	10	15%
Heterosexual contact	2	100%	4	100%	-	-	-	-	2	67%	53	79%
Other/unknown	-	-	-	-	-	-	-	-	1	33%	4	6%
Subtotal	2	100%	4	100%	1	100%	1	100%	3	100%	67	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	3	75%
Other/unknown	-	-	-	-	-	-	-	-	-	-	1	25%
Subtotal	-	-	-	-	-	-	-	-	-	-	4	100%
Total	13		12		12		11		8		410	

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	23%	5	42%	7	58%	7	64%	3	38%	208	51%
Injection drug use (IDU) only	-	-	-	-	-	-	1	9%	-	-	12	3%
Heterosexual contact only	3	23%	4	33%	1	8%	1	9%	3	38%	74	18%
Male-to-male sexual contact & IDU	-	-	-	-	1	8%	-	-	-	-	14	3%
IDU & Heterosexual contact	-	-	-	-	1	8%	-	-	-	-	13	3%
Male-to-male sexual contact & Heterosexual contact	1	8%	1	8%	-	-	1	9%	-	-	19	5%
Male-to-male sexual contact & IDU & Heterosexual contact	1	8%	-	-	-	-	-	-	-	-	3	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	3	1%
Other/unknown	5	38%	2	17%	2	17%	1	9%	2	25%	64	16%
Total	13		12		12		11		8		410	

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 10

Counties: Allen, Auglaize, Champaign, Hancock, Hardin, Logan, Mercer, Paulding, Putnam, Shelby, Van Wert

As of June 30, 2019

Reported persons living with diagnosed HIV infection in 2018 by current disease status and selected characteristics, HIV Prevention Planning Region 10

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2018			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	117.7	300	74%	127	70%	173	78%
Females	40.3	103	26%	55	30%	48	22%
Age at end of year							
<13	*	1	<1%	1	1%	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	28.0	9	2%	7	4%	2	1%
25-29	75.0	24	6%	15	8%	9	4%
30-34	125.0	36	9%	22	12%	14	6%
35-39	120.5	36	9%	21	12%	15	7%
40-44	161.6	46	11%	22	12%	24	11%
45-49	172.3	53	13%	20	11%	33	15%
50-54	167.3	55	14%	19	10%	36	16%
55-64	156.9	113	28%	43	24%	70	32%
65+	33.3	30	7%	12	7%	18	8%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	4	1%	1	1%	3	1%
Black/African-American	467.0	84	21%	39	21%	45	20%
Hispanic/Latinx	176.6	28	7%	13	7%	15	7%
White	58.9	272	67%	122	67%	150	68%
Multi-Race	166.1	15	4%	7	4%	8	4%
Unknown	*	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	3	1%	1	1%	2	1%
Asian/Pacific Islander Females	*	1	<1%	-	-	1	<1%
Black/African-American Males	488.5	47	12%	21	12%	26	12%
Black/African-American Females	442.2	37	9%	18	10%	19	9%
Hispanic/Latino Males	236.7	20	5%	8	4%	12	5%
Hispanic/Latina Females	108.0	8	2%	5	3%	3	1%
White Males	94.5	217	54%	91	50%	126	57%
White Females	23.7	55	14%	31	17%	24	11%
Multi-Race Males	287.3	13	3%	6	3%	7	3%
Multi-Race Females	*	2	<1%	1	1%	1	<1%
Unknown	*	-	-	-	-	-	-
Total	79.0	403		182		221	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by current disease status and transmission category, HIV Prevention Planning Region 10

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	196	66%	79	62%	117	68%
Injection drug use (IDU)	16	5%	8	6%	8	5%
Male-to-male sexual contact and IDU	17	6%	7	6%	10	6%
Heterosexual contact	27	9%	10	8%	17	10%
Other/unknown	43	14%	23	18%	20	12%
Subtotal	299	100%	127	100%	172	100%
Female adult or adolescent						
Injection drug use	15	15%	8	15%	7	15%
Heterosexual contact	77	77%	39	74%	38	81%
Other/unknown	8	8%	6	11%	2	4%
Subtotal	100	100%	53	100%	47	100%
Child (<13 yrs at diagnosis)						
Perinatal	4	100%	2	100%	2	100%
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	2	100%	2	100%
Total	403		182		221	

Reported persons living with diagnosed HIV infection in 2018 by current disease status and exposure category, HIV Prevention Planning Region 10

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	175	43%	71	39%	104	47%
Injection drug use (IDU) only	9	2%	4	2%	5	2%
Heterosexual contact only	105	26%	49	27%	56	25%
Male-to-male sexual contact & IDU	13	3%	7	4%	6	3%
IDU & Heterosexual contact	22	5%	12	7%	10	5%
Male-to-male sexual contact & Heterosexual contact	21	5%	8	4%	13	6%
Male-to-male sexual contact & IDU & Heterosexual contact	4	1%	-	-	4	2%
Perinatal exposure	4	1%	2	1%	2	1%
Other/unknown	50	12%	29	16%	21	10%
Total	403		182		221	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and age, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2018

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	-	-	*	1	4%	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	3	4%	*	1	4%	*	4	1%	*	1	7%	-	-
25-29	*	-	-	*	-	-	523.2	8	10%	*	4	14%	42.6	12	4%	*	-	-	-	-
30-34	*	-	-	*	-	-	1,044.4	12	14%	*	3	11%	81.9	21	8%	*	-	-	-	-
35-39	*	-	-	*	-	-	563.2	7	8%	*	2	7%	89.8	24	9%	*	3	20%	-	-
40-44	*	-	-	*	1	25%	983.3	10	12%	648.6	6	21%	112.1	29	11%	*	-	-	-	-
45-49	*	-	-	*	1	25%	865.4	9	11%	*	3	11%	127.4	36	13%	*	4	27%	-	-
50-54	*	-	-	*	-	-	1,235.7	14	17%	801.3	5	18%	111.4	34	13%	*	2	13%	-	-
55-64	*	-	-	*	2	50%	838.4	18	21%	*	2	7%	128.7	87	32%	*	4	27%	-	-
65+	*	-	-	*	-	-	*	3	4%	*	1	4%	29.1	25	9%	*	1	7%	-	-
Total	*	-	-	*	4	-	467.0	84	-	176.6	28	-	58.9	272	-	166.1	15	-	-	-

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2018

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	27	57%	7	37%	151	70%	11	85%	-	-
Injection drug use (IDU)	-	-	-	-	4	9%	2	11%	10	5%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	-	-	2	4%	3	16%	12	6%	-	-	-	-
Heterosexual contact	-	-	2	67%	8	17%	6	32%	11	5%	-	-	-	-
Other/unknown	-	-	1	33%	6	13%	1	5%	33	15%	2	15%	-	-
Subtotal	-	-	3	100%	47	100%	19	100%	217	100%	13	100%	-	-
Female adult or adolescent														
Injection drug use	-	-	-	-	4	11%	-	-	10	19%	1	50%	-	-
Heterosexual contact	-	-	1	100%	30	81%	6	86%	39	74%	1	50%	-	-
Other/unknown	-	-	-	-	3	8%	1	14%	4	8%	-	-	-	-
Subtotal	-	-	1	100%	37	100%	7	100%	53	100%	2	100%	-	-
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	-	-	2	100%	2	100%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	2	100%	2	100%	-	-	-	-
Total	-	-	4	-	84	-	28	-	272	-	15	-	-	-

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 10

Living with diagnosed HIV infection in 2018

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	24	29%	7	25%	136	50%	8	53%	-	-
Injection drug use (IDU) only	-	-	-	-	4	5%	-	-	5	2%	-	-	-	-
Heterosexual contact only	-	-	3	75%	38	45%	12	43%	51	19%	1	7%	-	-
Male-to-male sexual contact & IDU	-	-	-	-	2	2%	3	11%	8	3%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	4	5%	2	7%	15	6%	1	7%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	3	4%	-	-	15	6%	3	20%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	4	1%	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	2	7%	2	1%	-	-	-	-
Other/unknown	-	-	1	25%	9	11%	2	7%	36	13%	2	13%	-	-
Total	-	-	4	-	84	-	28	-	272	-	15	-	-	-

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 10, 2014-2018

Living with Diagnosed HIV Infection															
Characteristic	2014			2015			2016			2017			2018		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	100.5	257	76%	109.5	280	77%	109.4	279	74%	113.7	290	74%	117.7	300	74%
Females	32.1	83	24%	33.4	86	23%	38.2	98	26%	40.3	103	26%	40.3	103	26%
Age at end of year															
<13	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	2	1%	*	2	1%	*	1	<1%	*	1	<1%	*	-	-
20-24	*	4	1%	23.3	8	2%	23.9	8	2%	30.6	10	3%	28.0	9	2%
25-29	86.5	25	7%	102.4	30	8%	95.8	29	8%	82.7	26	7%	75.0	24	6%
30-34	81.2	24	7%	98.5	29	8%	106.0	31	8%	115.0	33	8%	125.0	36	9%
35-39	127.4	37	11%	140.2	41	11%	142.2	42	11%	123.5	37	9%	120.5	36	9%
40-44	158.1	49	14%	129.1	39	11%	124.4	36	10%	157.8	45	11%	161.6	46	11%
45-49	147.4	48	14%	163.2	52	14%	168.3	53	14%	166.2	52	13%	172.3	53	13%
50-54	164.6	62	18%	172.1	63	17%	185.5	66	18%	195.5	67	17%	167.3	55	14%
55-64	97.5	69	20%	113.2	81	22%	125.4	90	24%	134.6	97	25%	156.9	113	28%
65+	22.9	19	6%	23.6	20	5%	23.2	20	5%	27.3	24	6%	33.3	30	7%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	1	<1%	*	1	<1%	*	3	1%	*	2	1%	*	4	1%
Black/African-American	384.8	69	20%	392.9	71	19%	442.9	79	21%	435.2	78	20%	467.0	84	21%
Hispanic/Latinx	105.3	15	4%	128.7	19	5%	126.7	19	5%	163.8	25	6%	176.6	28	7%
White	52.1	244	72%	55.9	261	71%	56.6	263	70%	59.2	274	70%	58.9	272	67%
Multi-Race	135.6	11	3%	167.6	14	4%	150.4	13	3%	158.2	14	4%	166.1	15	4%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	<1%	*	1	<1%	*	2	1%	*	1	<1%	*	3	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	1	<1%	*	1	<1%	*	1	<1%
Black/African-American Males	425.2	41	12%	442.7	43	12%	493.5	47	12%	480.6	46	12%	488.5	47	12%
Black/African-American Females	337.8	28	8%	334.9	28	8%	384.8	32	8%	383.2	32	8%	442.2	37	9%
Hispanic/Latino Males	133.1	10	3%	167.2	13	4%	177.4	14	4%	211.2	17	4%	236.7	20	5%
Hispanic/Latina Females	74.2	5	1%	85.8	6	2%	70.4	5	1%	110.9	8	2%	108.0	8	2%
White Males	84.4	196	58%	91.6	212	58%	89.3	206	55%	93.4	215	55%	94.5	217	54%
White Females	20.3	48	14%	20.8	49	13%	24.4	57	15%	25.3	59	15%	23.7	55	14%
Multi-Race Males	222.9	9	3%	264.4	11	3%	231.6	10	3%	247.5	11	3%	287.3	13	3%
Multi-Race Females	*	2	1%	*	3	1%	*	3	1%	*	3	1%	*	2	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	66.1	340		71.3	366		73.7	377		76.9	393		79.0	403	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 10, 2014-2018

Transmission Category ^a	Living with Diagnosed HIV Infection									
	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	165	64%	186	67%	188	68%	199	69%	196	66%
Injection drug use (IDU)	11	4%	12	4%	15	5%	15	5%	16	5%
Male-to-male sexual contact and IDU	17	7%	17	6%	13	5%	11	4%	17	6%
Heterosexual contact	20	8%	20	7%	21	8%	24	8%	27	9%
Other/unknown	43	17%	44	16%	41	15%	40	14%	43	14%
Subtotal	256	100%	279	100%	278	100%	289	100%	299	100%
Female adult or adolescent										
Injection drug use	9	11%	11	13%	16	17%	18	18%	15	15%
Heterosexual contact	67	83%	68	81%	75	78%	76	75%	77	77%
Other/unknown	5	6%	5	6%	5	5%	7	7%	8	8%
Subtotal	81	100%	84	100%	96	100%	101	100%	100	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	100%	3	100%	3	100%	3	100%	4	100%
Other/unknown	-	-	-	-	-	-	-	-	-	-
Subtotal	3	100%	3	100%	3	100%	3	100%	4	100%
Total	340		366		377		393		403	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 10, 2014-2018

Exposure Category ^b	Living with Diagnosed HIV Infection									
	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	148	44%	167	46%	169	45%	181	46%	175	43%
Injection drug use (IDU) only	8	2%	9	2%	7	2%	10	3%	9	2%
Heterosexual contact only	88	26%	89	24%	97	26%	101	26%	105	26%
Male-to-male sexual contact & IDU	11	3%	11	3%	9	2%	7	2%	13	3%
IDU & Heterosexual contact	12	4%	14	4%	24	6%	23	6%	22	5%
Male-to-male sexual contact & Heterosexual contact	17	5%	19	5%	19	5%	18	5%	21	5%
Male-to-male sexual contact & IDU & Heterosexual contact	6	2%	6	2%	4	1%	4	1%	4	1%
Perinatal exposure	3	1%	3	1%	3	1%	3	1%	4	1%
Other/unknown	47	14%	48	13%	45	12%	46	12%	50	12%
Total	340		366		377		393		403	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.