

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan, Muskingum, Noble, Perry, Washington

As of June 30, 2019

Reported new diagnoses of HIV infection in 2018 by disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	2018 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	2.4	5	63%	2	40%	2	100%	1	100%
Females	*	3	38%	3	60%	-	-	-	-
Age at diagnosis (yr)									
<13	*	-	-	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-	-	-
20-24	*	1	13%	1	20%	-	-	-	-
25-29	*	-	-	-	-	-	-	-	-
30-34	*	3	38%	3	60%	-	-	-	-
35-39	*	1	13%	-	-	1	50%	-	-
40-44	*	-	-	-	-	-	-	-	-
45-49	*	3	38%	1	20%	1	50%	1	100%
50-54	*	-	-	-	-	-	-	-	-
55-64	*	-	-	-	-	-	-	-	-
65+	*	-	-	-	-	-	-	-	-
Race/Ethnicity ^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-	-	-
Black/African-American	*	1	13%	1	20%	-	-	-	-
Hispanic/Latinx	*	-	-	-	-	-	-	-	-
White	1.8	7	88%	4	80%	2	100%	1	100%
Multi-Race	*	-	-	-	-	-	-	-	-
Race/Ethnicity ^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-	-	-
Black/African-American Males	*	-	-	-	-	-	-	-	-
Black/African-American Females	*	1	13%	1	20%	-	-	-	-
Hispanic/Latino Males	*	-	-	-	-	-	-	-	-
Hispanic/Latina Females	*	-	-	-	-	-	-	-	-
White Males	2.6	5	63%	2	40%	2	100%	1	100%
White Females	*	2	25%	2	40%	-	-	-	-
Multi-Race Males	*	-	-	-	-	-	-	-	-
Multi-Race Females	*	-	-	-	-	-	-	-	-
Total	1.9	8		5		2		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	3	60%	2	100%	-	-	1	100%
Injection drug use (IDU)	1	20%	-	-	1	50%	-	-
Male-to-male sexual contact and IDU	1	20%	-	-	1	50%	-	-
Heterosexual contact	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	5	100%	2	100%	2	100%	1	100%
Female adult or adolescent								
Injection drug use	1	33%	1	33%	-	-	-	-
Heterosexual contact	1	33%	1	33%	-	-	-	-
Other/unknown	1	33%	1	33%	-	-	-	-
Subtotal	3	100%	3	100%	-	-	-	-
Child (<13 yrs at diagnosis)								
Perinatal	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-
Total	8		5		2		1	

Reported new diagnoses of HIV infection in 2018 by disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3	38%	2	40%	-	-	1	100%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-
Heterosexual contact only	1	13%	1	20%	-	-	-	-
Male-to-male sexual contact & IDU	1	13%	-	-	1	50%	-	-
IDU & Heterosexual contact	2	25%	1	20%	1	50%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-
Other/unknown	1	13%	1	20%	-	-	-	-
Total	8		5		2		1	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 6

2018 diagnosis of HIV infection																		
Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx*			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	*	-	-	*	-	-	*	-	-	*	-	-	*	1	14%	*	-	-
25-29	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	*	3	43%	*	-	-
35-39	*	-	-	*	-	-	*	-	-	*	-	-	*	1	14%	*	-	-
40-44	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
45-49	*	-	-	*	-	-	*	1	100%	*	-	-	*	2	29%	*	-	-
50-54	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
55-64	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
65+	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Total	*	-	-	*	-	-	*	1	-	*	-	-	1.8	7	-	*	-	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

2018 diagnosis of HIV infection												
Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent	-	-	-	-	-	-	-	-	3	60%	-	-
Male-to-male sexual contact	-	-	-	-	-	-	-	-	1	20%	-	-
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	20%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	5	100%	-	-
Female adult or adolescent	-	-	-	-	-	-	-	-	1	50%	-	-
Injection drug use	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	1	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-
Subtotal	-	-	-	-	1	100%	-	-	2	100%	-	-
Child (<13 yrs at diagnosis)	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	1	-	-	-	7	-	-	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

2018 diagnosis of HIV infection												
Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	-	-	-	-	3	43%	-	-
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	1	100%	-	-	-	-	-	-
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	14%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	2	29%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	14%	-	-
Total	-	-	-	-	1	-	-	-	7	-	-	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Diagnosis of HIV Infection by Year															Cumulative Diagnoses	
	2014			2015			2016			2017			2018				
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	5.2	11	100%	2.8	6	86%	2.8	6	67%	4.7	10	71%	2.4	5	63%	376	81%
Females	*	-	-	*	1	14%	*	3	33%	*	4	29%	*	3	38%	89	19%
Age at diagnosis (yr)																	
<13	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	6	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	7	2%
20-24	*	2	18%	*	1	14%	*	4	44%	*	4	29%	*	1	13%	51	11%
25-29	*	1	9%	*	1	14%	*	1	11%	19.1	5	36%	*	-	-	71	15%
30-34	*	1	9%	*	2	29%	*	1	11%	*	-	-	*	3	38%	79	17%
35-39	*	-	-	*	-	-	*	-	-	*	3	21%	*	1	13%	83	18%
40-44	*	4	36%	*	1	14%	*	-	-	*	1	7%	*	-	-	65	14%
45-49	*	-	-	*	1	14%	*	2	22%	*	1	7%	*	3	38%	45	10%
50-54	*	1	9%	*	-	-	*	-	-	*	-	-	*	-	-	25	5%
55-64	*	1	9%	*	1	14%	*	1	11%	*	-	-	*	-	-	23	5%
65+	*	1	9%	*	-	-	*	-	-	*	-	-	*	-	-	8	2%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American	*	-	-	*	1	14%	*	-	-	*	1	7%	*	1	13%	36	8%
Hispanic/Latinx	*	2	18%	*	-	-	*	1	11%	*	2	14%	*	-	-	15	3%
White	2.3	9	82%	1.5	6	86%	2.0	8	89%	2.5	10	71%	1.8	7	88%	402	86%
Multi-Race	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-	8	2%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Race/Ethnicity ^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Black/African-American Males	*	-	-	*	1	14%	*	-	-	*	-	-	*	-	-	21	5%
Black/African-American Females	*	-	-	*	-	-	*	-	-	*	1	7%	*	1	13%	15	3%
Hispanic/Latino Males	*	2	18%	*	-	-	*	1	11%	*	2	14%	*	-	-	12	3%
Hispanic/Latina Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	3	1%
White Males	4.6	9	82%	2.5	5	71%	2.5	5	56%	3.6	7	50%	2.6	5	63%	333	72%
White Females	*	-	-	*	1	14%	*	3	33%	*	3	21%	*	2	25%	69	15%
Multi-Race Males	*	-	-	*	-	-	*	-	-	*	1	7%	*	-	-	8	2%
Multi-Race Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	4	1%
Total	2.6	11		1.6	7		2.1	9		3.3	14		1.9	8		465	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Diagnosis of HIV Infection by Year										Cumulative Diagnoses	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	6	55%	3	50%	4	67%	6	60%	3	60%	230	62%
Injection drug use (IDU)	-	-	-	-	-	-	1	10%	1	20%	13	4%
Male-to-male sexual contact and IDU	1	9%	-	-	-	-	2	20%	1	20%	18	5%
Heterosexual contact	1	9%	-	-	-	-	1	10%	-	-	25	7%
Other/unknown	3	27%	3	50%	2	33%	-	-	-	-	85	23%
Subtotal	11	100%	6	100%	6	100%	10	100%	5	100%	371	100%
Female adult or adolescent												
Injection drug use	-	-	1	100%	1	33%	3	75%	1	33%	15	17%
Heterosexual contact	-	-	-	-	2	67%	1	25%	1	33%	60	68%
Other/unknown	-	-	-	-	-	-	-	-	1	33%	13	15%
Subtotal	-	-	1	100%	3	100%	4	100%	3	100%	88	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	4	67%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	33%
Subtotal	-	-	-	-	-	-	-	-	-	-	6	100%
Total	11		7		9		14		8		465	

Trends in reported diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Diagnosis of HIV Infection by Year										Cumulative Diagnoses	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	6	55%	2	29%	4	44%	5	36%	3	38%	216	46%
Injection drug use (IDU) only	-	-	-	-	-	-	1	7%	-	-	12	3%
Heterosexual contact only	1	9%	-	-	2	22%	2	14%	1	13%	85	18%
Male-to-male sexual contact & IDU	1	9%	-	-	-	-	1	7%	1	13%	14	3%
IDU & Heterosexual contact	-	-	1	14%	1	11%	3	21%	2	25%	16	3%
Male-to-male sexual contact & Heterosexual contact	-	-	1	14%	-	-	1	7%	-	-	14	3%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	1	7%	-	-	4	1%
Perinatal exposure	-	-	-	-	-	-	-	-	-	-	5	1%
Other/unknown	3	27%	3	43%	2	22%	-	-	1	13%	99	21%
Total	11		7		9		14		8		465	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 6

Characteristic	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	4	100%	1	100%	3	100%
Females	-	-	-	-	-	-
Age at AIDS diagnosis (yr)						
<13	-	-	-	-	-	-
13-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	-	-	-	-	-	-
35-39	1	25%	-	-	1	33%
40-44	-	-	-	-	-	-
45-49	2	50%	-	-	2	67%
50-54	1	25%	1	100%	-	-
55-64	-	-	-	-	-	-
65+	-	-	-	-	-	-
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African-American	-	-	-	-	-	-
Hispanic/Latinx	-	-	-	-	-	-
White	4	100%	1	100%	3	100%
Multi-Race	-	-	-	-	-	-
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	-	-	-	-	-	-
Black/African-American Females	-	-	-	-	-	-
Hispanic/Latino Males	-	-	-	-	-	-
Hispanic/Latina Females	-	-	-	-	-	-
White Males	4	100%	1	100%	3	100%
White Females	-	-	-	-	-	-
Multi-Race Males	-	-	-	-	-	-
Multi-Race Females	-	-	-	-	-	-
Total	4		1		3	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 6

Transmission Category ^b	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	2	50%	1	100%	1	33%
Injection drug use (IDU)	1	25%	-	-	1	33%
Male-to-male sexual contact and IDU	1	25%	-	-	1	33%
Heterosexual contact	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	4	100%	1	100%	3	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
Total	4		1		3	

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 6

Exposure Category ^c	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2	50%	1	100%	1	33%
Injection drug use (IDU) only	-	-	-	-	-	-
Heterosexual contact only	-	-	-	-	-	-
Male-to-male sexual contact & IDU	1	25%	-	-	1	33%
IDU & Heterosexual contact	1	25%	-	-	1	33%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-
Total	4		1		3	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 6

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex at birth												
Males	5	100%	3	100%	3	100%	6	86%	4	100%	251	87%
Females	-	-	-	-	-	-	1	14%	-	-	38	13%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	-	-	2	1%
13-14	-	-	-	-	-	-	-	-	-	-	2	1%
15-19	-	-	-	-	-	-	-	-	-	-	2	1%
20-24	-	-	-	-	-	-	-	-	-	-	5	2%
25-29	-	-	-	-	-	-	2	29%	-	-	32	11%
30-34	-	-	-	-	1	33%	-	-	-	-	45	16%
35-39	-	-	-	-	-	-	2	29%	1	25%	55	19%
40-44	1	20%	-	-	-	-	-	-	-	-	60	21%
45-49	-	-	1	33%	-	-	1	14%	2	50%	35	12%
50-54	2	40%	1	33%	2	67%	-	-	1	25%	25	9%
55-64	1	20%	1	33%	-	-	1	14%	-	-	22	8%
65+	1	20%	-	-	-	-	1	14%	-	-	4	1%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American	-	-	-	-	-	-	1	14%	-	-	19	7%
Hispanic/Latinx	-	-	-	-	-	-	1	14%	-	-	5	2%
White	5	100%	3	100%	3	100%	4	57%	4	100%	259	90%
Multi-Race	-	-	-	-	-	-	1	14%	-	-	6	2%
Race/Ethnicity ^a and Sex at birth												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	-	-
Black/African-American Males	-	-	-	-	-	-	-	-	-	-	14	5%
Black/African-American Females	-	-	-	-	-	-	1	14%	-	-	5	2%
Hispanic/Latino Males	-	-	-	-	-	-	1	14%	-	-	4	1%
Hispanic/Latina Females	-	-	-	-	-	-	-	-	-	-	1	<1%
White Males	5	100%	3	100%	3	100%	4	57%	4	100%	227	79%
White Females	-	-	-	-	-	-	-	-	-	-	32	11%
Multi-Race Males	-	-	-	-	-	-	1	14%	-	-	6	2%
Multi-Race Females	-	-	-	-	-	-	-	-	-	-	-	-
Total	5		3		3		7		4		289	

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	1	20%	1	33%	2	67%	5	83%	2	50%	151	61%
Injection drug use (IDU)	-	-	-	-	-	-	-	-	1	25%	12	5%
Male-to-male sexual contact and IDU	-	-	-	-	-	-	-	-	1	25%	15	6%
Heterosexual contact	-	-	-	-	-	-	1	17%	-	-	18	7%
Other/unknown	4	80%	2	67%	1	33%	-	-	-	-	51	21%
Subtotal	5	100%	3	100%	3	100%	6	100%	4	100%	247	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	-	-	1	100%	-	-	7	18%
Heterosexual contact	-	-	-	-	-	-	-	-	-	-	27	71%
Other/unknown	-	-	-	-	-	-	-	-	-	-	4	11%
Subtotal	-	-	-	-	-	-	1	100%	-	-	38	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	2	50%
Other/unknown	-	-	-	-	-	-	-	-	-	-	2	50%
Subtotal	-	-	-	-	-	-	-	-	-	-	4	100%
Total	5		3		3		7		4		289	

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	1	20%	1	33%	2	67%	5	71%	2	50%	144	50%
Injection drug use (IDU) only	-	-	-	-	-	-	-	-	-	-	9	3%
Heterosexual contact only	-	-	-	-	-	-	1	14%	-	-	45	16%
Male-to-male sexual contact & IDU	-	-	-	-	-	-	-	-	1	25%	12	4%
IDU & Heterosexual contact	-	-	-	-	-	-	1	14%	1	25%	10	3%
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	7	2%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	3	1%
Perinatal exposure	-	-	-	-	1	33%	-	-	-	-	3	1%
Other/unknown	4	80%	2	67%	-	-	-	-	-	-	56	19%
Total	5		3		3		7		4		289	

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 6

Counties: Athens, Belmont, Guernsey, Meigs, Monroe, Morgan, Muskingum, Noble, Perry, Washington

As of June 30, 2019

Reported persons living with diagnosed HIV infection in 2018 by current disease status and selected characteristics, HIV Prevention Planning Region 6

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2018			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	127.1	267	79%	122	76%	145	82%
Females	33.3	70	21%	38	24%	32	18%
Age at end of year							
<13	*	2	1%	1	1%	1	1%
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	21.6	7	2%	5	3%	2	1%
25-29	91.2	24	7%	19	12%	5	3%
30-34	89.3	21	6%	18	11%	3	2%
35-39	111.0	27	8%	17	11%	10	6%
40-44	97.3	22	7%	11	7%	11	6%
45-49	174.3	45	13%	27	17%	18	10%
50-54	226.8	61	18%	17	11%	44	25%
55-64	167.3	102	30%	38	24%	64	36%
65+	32.8	26	8%	7	4%	19	11%
Race/Ethnicity^b							
American Indian/Alaska Native	*	-	-	-	-	-	-
Asian/Pacific Islander	*	2	1%	1	1%	1	1%
Black/African-American	257.0	27	8%	9	6%	18	10%
Hispanic/Latinx	301.1	15	4%	8	5%	7	4%
White	71.0	278	82%	136	85%	142	80%
Multi-Race	132.8	11	3%	2	1%	9	5%
Unknown	*	4	1%	4	3%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-
Asian/Pacific Islander Males	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Females	*	-	-	-	-	-	-
Black/African-American Males	288.0	19	6%	5	3%	14	8%
Black/African-American Females	204.7	8	2%	4	3%	4	2%
Hispanic/Latino Males	472.6	12	4%	7	4%	5	3%
Hispanic/Latina Females	*	3	1%	1	1%	2	1%
White Males	114.7	223	66%	105	66%	118	67%
White Females	27.9	55	16%	31	19%	24	14%
Multi-Race Males	217.2	9	3%	2	1%	7	4%
Multi-Race Females	*	2	1%	-	-	2	1%
Unknown	*	4	1%	4	3%	-	-
Total	80.1	337		160		177	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by current disease status and transmission category, HIV Prevention Planning Region 6

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	187	71%	87	71%	100	70%
Injection drug use (IDU)	8	3%	3	2%	5	4%
Male-to-male sexual contact and IDU	18	7%	9	7%	9	6%
Heterosexual contact	14	5%	2	2%	12	8%
Other/unknown	37	14%	21	17%	16	11%
Subtotal	264	100%	122	100%	142	100%
Female adult or adolescent						
Injection drug use	12	17%	6	16%	6	19%
Heterosexual contact	51	74%	25	68%	26	81%
Other/unknown	6	9%	6	16%	-	-
Subtotal	69	100%	37	100%	32	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	75%	1	100%	2	67%
Other/unknown	1	25%	-	-	1	33%
Subtotal	4	100%	1	100%	3	100%
Total	337		160		177	

Reported persons living with diagnosed HIV infection in 2018 by current disease status and exposure category, HIV Prevention Planning Region 6

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	179	53%	84	53%	95	54%
Injection drug use (IDU) only	10	3%	3	2%	7	4%
Heterosexual contact only	65	19%	27	17%	38	21%
Male-to-male sexual contact & IDU	15	4%	8	5%	7	4%
IDU & Heterosexual contact	10	3%	6	4%	4	2%
Male-to-male sexual contact & Heterosexual contact	8	2%	3	2%	5	3%
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	1	1%	2	1%
Perinatal exposure	3	1%	1	1%	2	1%
Other/unknown	44	13%	27	17%	17	10%
Total	337		160		177	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and age, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2018

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	-	-	*	1	4%	*	1	7%	*	-	-	*	-	-	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	-	-
20-24	*	-	-	*	-	-	*	1	4%	*	-	-	21.3	6	2%	*	-	-	-	-
25-29	*	-	-	*	-	-	*	1	4%	*	3	20%	80.2	19	7%	*	1	9%	-	-
30-34	*	-	-	*	-	-	*	-	-	*	-	-	96.6	21	8%	*	-	-	-	-
35-39	*	-	-	*	-	-	*	3	11%	*	2	13%	97.1	22	8%	*	-	-	-	-
40-44	*	-	-	*	-	-	*	1	4%	*	-	-	89.7	19	7%	*	2	18%	-	-
45-49	*	-	-	*	1	50%	1,071.4	6	22%	*	2	13%	142.7	35	13%	*	-	-	1	25%
50-54	*	-	-	*	1	50%	*	4	15%	*	3	20%	191.0	49	18%	*	3	27%	1	25%
55-64	*	-	-	*	-	-	803.9	10	37%	*	3	20%	140.4	82	29%	854.7	5	45%	2	50%
65+	*	-	-	*	-	-	*	-	-	*	1	7%	32.7	25	9%	*	-	-	-	-
Total	*	-	-	*	2	-	257.0	27	-	301.1	15	-	71.0	278	-	132.8	11	-	4	-

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2018

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	-	-	-	-	9	50%	6	50%	163	74%	9	100%	-	-
Injection drug use (IDU)	-	-	-	-	2	11%	-	-	6	3%	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	50%	-	-	1	8%	16	7%	-	-	-	-
Heterosexual contact	-	-	-	-	1	6%	3	25%	10	5%	-	-	-	-
Other/unknown	-	-	1	50%	6	33%	2	17%	26	12%	-	-	2	100%
Subtotal	-	-	2	100%	18	100%	12	100%	221	100%	9	100%	2	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	1	13%	-	-	10	18%	1	50%	-	-
Heterosexual contact	-	-	-	-	6	75%	2	100%	42	76%	1	50%	-	-
Other/unknown	-	-	-	-	1	13%	-	-	3	5%	-	-	2	100%
Subtotal	-	-	-	-	8	100%	2	100%	55	100%	2	100%	2	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	-	-	1	100%	1	100%	1	50%	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	1	50%	-	-	-	-
Subtotal	-	-	-	-	1	100%	1	100%	2	100%	-	-	-	-
Total	-	-	2	-	27	-	15	-	278	-	11	-	4	-

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 6

Living with diagnosed HIV infection in 2018

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	8	30%	6	40%	156	56%	9	82%	-	-
Injection drug use (IDU) only	-	-	-	-	2	7%	-	-	7	3%	1	9%	-	-
Heterosexual contact only	-	-	-	-	7	26%	5	33%	52	19%	1	9%	-	-
Male-to-male sexual contact & IDU	-	-	1	50%	-	-	-	-	14	5%	-	-	-	-
IDU & Heterosexual contact	-	-	-	-	1	4%	-	-	9	3%	-	-	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	1	4%	-	-	7	3%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	1	7%	2	1%	-	-	-	-
Perinatal exposure	-	-	-	-	1	4%	1	7%	1	0%	-	-	-	-
Other/unknown	-	-	1	50%	7	26%	2	13%	30	11%	-	-	4	100%
Total	-	-	2	-	27	-	15	-	278	-	11	-	4	-

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 6, 2014-2018

Characteristic	Living with Diagnosed HIV Infection														
	2014			2015			2016			2017			2018		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	110.3	233	81%	109.4	232	81%	117.3	248	81%	117.1	247	80%	127.1	267	79%
Females	26.4	56	19%	26.4	56	19%	26.9	57	19%	28.8	61	20%	33.3	70	21%
Age at end of year															
<13	*	2	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
20-24	20.1	7	2%	*	4	1%	17.8	6	2%	18.0	6	2%	21.6	7	2%
25-29	40.3	10	3%	54.6	14	5%	65.4	17	6%	84.1	22	7%	91.2	24	7%
30-34	83.1	20	7%	78.9	19	7%	92.4	22	7%	67.8	16	5%	89.3	21	6%
35-39	61.2	14	5%	81.0	19	7%	101.0	24	8%	112.1	27	9%	111.0	27	8%
40-44	150.4	38	13%	110.2	27	9%	106.6	25	8%	104.9	24	8%	97.3	22	7%
45-49	192.2	51	18%	208.7	55	19%	170.8	45	15%	175.9	46	15%	174.3	45	13%
50-54	187.8	57	20%	188.9	56	19%	216.9	62	20%	202.7	56	18%	226.8	61	18%
55-64	122.4	74	26%	125.8	77	27%	138.6	85	28%	143.1	88	29%	167.3	102	30%
65+	22.1	16	6%	20.3	15	5%	22.5	17	6%	27.1	21	7%	32.8	26	8%
Race/Ethnicity^b															
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander	*	1	<1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
Black/African-American	216.6	23	8%	215.0	23	8%	249.8	26	9%	228.6	24	8%	257.0	27	8%
Hispanic/Latinx	160.5	7	2%	173.7	8	3%	253.6	12	4%	267.5	13	4%	301.1	15	4%
White	61.3	243	84%	60.8	241	84%	63.3	250	82%	64.7	255	83%	71.0	278	82%
Multi-Race	142.5	11	4%	126.8	10	3%	136.8	11	4%	123.1	10	3%	132.8	11	3%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Asian/Pacific Islander Males	*	1	<1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
Asian/Pacific Islander Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
Black/African-American Males	241.6	16	6%	255.3	17	6%	307.0	20	7%	274.1	18	6%	288.0	19	6%
Black/African-American Females	175.3	7	2%	148.7	6	2%	154.2	6	2%	152.7	6	2%	204.7	8	2%
Hispanic/Latino Males	225.2	5	2%	304.2	7	2%	421.2	10	3%	452.3	11	4%	472.6	12	4%
Hispanic/Latina Females	*	2	1%	*	1	<1%	*	2	1%	*	2	1%	*	3	1%
White Males	101.9	200	69%	99.6	196	68%	104.5	205	67%	105.3	206	67%	114.7	223	66%
White Females	21.5	43	15%	22.5	45	16%	22.6	45	15%	24.7	49	16%	27.9	55	16%
Multi-Race Males	236.2	9	3%	204.1	8	3%	225.6	9	3%	198.1	8	3%	217.2	9	3%
Multi-Race Females	*	2	1%	*	2	1%	*	2	1%	*	2	1%	*	2	1%
Unknown	*	4	1%	*	4	1%	*	4	1%	*	4	1%	*	4	1%
Total	68.2	289		67.9	288		72.1	305		72.9	308		80.1	337	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 6, 2014-2018

Living with Diagnosed HIV Infection										
Transmission Category ^a	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	164	71%	163	71%	173	71%	174	71%	187	71%
Injection drug use (IDU)	6	3%	5	2%	6	2%	8	3%	8	3%
Male-to-male sexual contact and IDU	9	4%	9	4%	14	6%	13	5%	18	7%
Heterosexual contact	12	5%	12	5%	13	5%	13	5%	14	5%
Other/unknown	39	17%	40	17%	39	16%	36	15%	37	14%
Subtotal	230	100%	229	100%	245	100%	244	100%	264	100%
Female adult or adolescent										
Injection drug use	6	11%	7	13%	8	14%	9	15%	12	17%
Heterosexual contact	43	78%	43	78%	43	77%	46	77%	51	74%
Other/unknown	6	11%	5	9%	5	9%	5	8%	6	9%
Subtotal	55	100%	55	100%	56	100%	60	100%	69	100%
Child (<13 yrs at diagnosis)										
Perinatal	3	75%	3	75%	3	75%	3	75%	3	75%
Other/unknown	1	25%	1	25%	1	25%	1	25%	1	25%
Subtotal	4	100%	4	100%	4	100%	4	100%	4	100%
Total	289		288		305		308		337	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 6, 2014-2018

Living with Diagnosed HIV Infection										
Exposure Category ^b	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	155	54%	153	53%	165	54%	165	54%	179	53%
Injection drug use (IDU) only	7	2%	7	2%	9	3%	8	3%	10	3%
Heterosexual contact only	55	19%	55	19%	56	18%	59	19%	65	19%
Male-to-male sexual contact & IDU	7	2%	6	2%	11	4%	9	3%	15	4%
IDU & Heterosexual contact	5	2%	5	2%	5	2%	9	3%	10	3%
Male-to-male sexual contact & Heterosexual contact	9	3%	10	3%	8	3%	9	3%	8	2%
Male-to-male sexual contact & IDU & Heterosexual contact	2	1%	3	1%	3	1%	4	1%	3	1%
Perinatal exposure	4	1%	4	1%	4	1%	3	1%	3	1%
Other/unknown	45	16%	45	16%	44	14%	42	14%	44	13%
Total	289		288		305		308		337	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.