

New Diagnoses of HIV Infection Reported in Franklin County

As of June 30, 2019

Reported new diagnoses of HIV infection in 2018 by disease status and selected characteristics, Franklin County

Characteristic	2018 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	23.7	152	75%	118	76%	23	66%	11	85%
Females	7.6	51	25%	37	24%	12	34%	2	15%
Age at diagnosis (yr)									
<13	*	4	2%	3	2%	-	-	1	8%
13-14	*	-	-	-	-	-	-	-	-
15-19	13.2	11	5%	9	6%	2	6%	-	-
20-24	30.7	28	14%	24	15%	2	6%	2	15%
25-29	41.0	52	26%	45	29%	6	17%	1	8%
30-34	31.8	36	18%	27	17%	7	20%	2	15%
35-39	19.9	19	9%	10	6%	6	17%	3	23%
40-44	18.7	15	7%	12	8%	2	6%	1	8%
45-49	20.1	16	8%	13	8%	3	9%	-	-
50-54	11.9	9	4%	4	3%	5	14%	-	-
55-64	6.0	9	4%	5	3%	2	6%	2	15%
65+	*	4	2%	3	2%	-	-	1	8%
Race/Ethnicity ^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	2	1%	2	1%	-	-	-	-
Black/African-American	39.0	117	58%	90	58%	21	60%	6	46%
Hispanic/Latinx	18.7	14	7%	10	6%	2	6%	2	15%
White	7.8	64	32%	49	32%	11	31%	4	31%
Multi-Race	15.5	6	3%	4	3%	1	3%	1	8%
Race/Ethnicity ^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	<1%	1	1%	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-	-	-
Black/African-American Males	60.9	87	43%	67	43%	14	40%	6	46%
Black/African-American Females	19.1	30	15%	23	15%	7	20%	-	-
Hispanic/Latino Males	30.7	12	6%	8	5%	2	6%	2	15%
Hispanic/Latina Females	*	2	1%	2	1%	-	-	-	-
White Males	12.2	49	24%	40	26%	7	20%	2	15%
White Females	3.6	15	7%	9	6%	4	11%	2	15%
Multi-Race Males	*	3	1%	2	1%	-	-	1	8%
Multi-Race Females	*	3	1%	2	1%	1	3%	-	-
Total	15.5	203		155		35		13	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by disease status and transmission category, Franklin County

Transmission Category ^a	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	102	68%	81	69%	15	65%	6	60%
Injection drug use (IDU)	6	4%	5	4%	1	4%	-	-
Male-to-male sexual contact and IDU	8	5%	8	7%	-	-	-	-
Heterosexual contact	4	3%	3	3%	-	-	1	10%
Other/unknown	30	20%	20	17%	7	30%	3	30%
Subtotal	150	100%	117	100%	23	100%	10	100%
Female adult or adolescent								
Injection drug use	11	22%	5	14%	5	42%	1	50%
Heterosexual contact	36	73%	29	83%	6	50%	1	50%
Other/unknown	2	4%	1	3%	1	8%	-	-
Subtotal	49	100%	35	100%	12	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	4	100%	3	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	4	100%	3	100%	-	-	1	100%
Total	203		155		35		13	

Reported new diagnoses of HIV infection in 2018 by disease status and exposure category, Franklin County

Exposure Category ^b	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	100	49%	79	51%	15	43%	6	46%
Injection drug use (IDU) only	8	4%	5	3%	3	9%	-	-
Heterosexual contact only	40	20%	32	21%	6	17%	2	15%
Male-to-male sexual contact & IDU	8	4%	8	5%	-	-	-	-
IDU & Heterosexual contact	9	4%	5	3%	3	9%	1	8%
Male-to-male sexual contact & Heterosexual contact	2	1%	2	1%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	4	2%	3	2%	-	-	1	8%
Other/unknown	32	16%	21	14%	8	23%	3	23%
Total	203		155		35		13	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and age at diagnosis, Franklin County

2018 diagnosis of HIV infection

Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	1	50%	*	3	3%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	50%	33.2	7	6%	*	-	-	*	3	5%	*	-	-
20-24	*	-	-	*	-	-	78.8	17	15%	*	4	29%	11.3	6	9%	*	1	17%
25-29	*	-	-	*	-	-	126.5	35	30%	*	2	14%	18.9	15	23%	*	-	-
30-34	*	-	-	*	-	-	64.7	15	13%	77.8	5	36%	19.5	14	22%	*	2	33%
35-39	*	-	-	*	-	-	52.4	11	9%	*	1	7%	10.1	6	9%	*	1	17%
40-44	*	-	-	*	-	-	38.2	7	6%	*	1	7%	14.1	7	11%	*	-	-
45-49	*	-	-	*	-	-	49.0	9	8%	*	1	7%	9.7	5	8%	*	1	17%
50-54	*	-	-	*	-	-	42.4	7	6%	*	-	-	*	2	3%	*	-	-
55-64	*	-	-	*	-	-	*	4	3%	*	-	-	4.7	5	8%	*	-	-
65+	*	-	-	*	-	-	*	2	2%	*	-	-	*	1	2%	*	1	17%
Total	*	-	-	*	2	-	39.0	117	-	18.7	14	-	7.8	64	-	15.5	6	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and transmission category, Franklin County

2018 diagnosis of HIV infection

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	54	64%	10	83%	36	73%	2	67%
Injection drug use (IDU)	-	-	-	-	1	1%	-	-	5	10%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	3	4%	-	-	5	10%	-	-
Heterosexual contact	-	-	-	-	2	2%	1	8%	1	2%	-	-
Other/unknown	-	-	1	100%	25	29%	1	8%	2	4%	1	33%
Subtotal	-	-	1	100%	85	100%	12	100%	49	100%	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	3%	-	-	10	67%	-	-
Heterosexual contact	-	-	-	-	27	93%	2	100%	5	33%	2	67%
Other/unknown	-	-	-	-	1	3%	-	-	-	-	1	33%
Subtotal	-	-	-	-	29	100%	2	100%	15	100%	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	3	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	3	100%	-	-	-	-	-	-
Total	-	-	2	-	117	-	14	-	64	-	6	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and exposure category, Franklin County

2018 diagnosis of HIV infection

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	54	46%	9	64%	35	55%	2	33%
Injection drug use (IDU) only	-	-	-	-	2	2%	-	-	6	9%	-	-
Heterosexual contact only	-	-	-	-	29	25%	3	21%	6	9%	2	33%
Male-to-male sexual contact & IDU	-	-	-	-	3	3%	-	-	5	8%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	9	14%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	1	7%	1	2%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	50%	3	3%	-	-	-	-	-	-
Other/unknown	-	-	1	50%	26	22%	1	7%	2	3%	2	33%
Total	-	-	2	-	117	-	14	-	64	-	6	-

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by selected characteristics, Franklin County

Diagnosis of HIV Infection by Year																	
Characteristic	2014			2015			2016			2017			2018			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	31.8	191	89%	24.1	147	73%	26.2	162	83%	27.9	176	79%	23.7	152	75%	6,309	84%
Females	3.8	24	11%	8.6	55	27%	5.3	34	17%	7.3	48	21%	7.6	51	25%	1,182	16%
Age at diagnosis (yr)																	
<13	*	1	<1%	*	2	1%	*	1	1%	*	-	-	*	4	2%	49	1%
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	8	<1%
15-19	7.6	6	3%	18.8	15	7%	13.7	11	6%	12.0	10	4%	13.2	11	5%	270	4%
20-24	49.6	46	21%	36.8	33	16%	39.1	35	18%	36.3	33	15%	30.7	28	14%	972	13%
25-29	40.5	48	22%	32.5	40	20%	28.7	36	18%	42.6	54	24%	41.0	52	26%	1,352	18%
30-34	35.1	35	16%	27.6	28	14%	22.1	23	12%	29.6	32	14%	31.8	36	18%	1,448	19%
35-39	28.5	24	11%	30.9	27	13%	28.9	26	13%	24.8	23	10%	19.9	19	9%	1,226	16%
40-44	21.2	17	8%	23.8	19	9%	24.3	19	10%	20.3	16	7%	18.7	15	7%	893	12%
45-49	26.1	20	9%	24.4	19	9%	15.1	12	6%	18.7	15	7%	20.1	16	8%	604	8%
50-54	8.6	7	3%	7.5	6	3%	20.6	16	8%	23.5	18	8%	11.9	9	4%	342	5%
55-64	6.5	9	4%	5.6	8	4%	11.1	16	8%	12.9	19	8%	6.0	9	4%	268	4%
65+	*	2	1%	3.6	5	2%	*	1	1%	*	4	2%	*	4	2%	59	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-	6	<1%
Asian/Pacific Islander	*	4	2%	11.7	7	3%	*	4	2%	*	3	1%	*	2	1%	48	1%
Black/African-American	38.2	102	47%	42.6	117	58%	37.9	107	55%	39.4	115	51%	39.0	117	58%	2,829	38%
Hispanic/Latinx	25.4	16	7%	20.0	13	6%	14.9	10	5%	18.3	13	6%	18.7	14	7%	293	4%
White	10.7	87	40%	7.4	60	30%	8.2	67	34%	10.4	85	38%	7.8	64	32%	4,076	54%
Multi-Race	17.9	6	3%	14.4	5	2%	22.1	8	4%	18.6	7	3%	15.5	6	3%	217	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	22	<1%
Race/Ethnicity ^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-	4	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	*	3	1%	*	4	2%	*	4	2%	*	1	<1%	*	1	<1%	34	<1%
Asian/Pacific Islander Females	*	1	<1%	*	3	1%	*	-	-	*	2	1%	*	1	<1%	14	<1%
Black/African-American Males	67.5	85	40%	57.8	75	37%	62.0	83	42%	59.9	83	37%	60.9	87	43%	2,061	28%
Black/African-American Females	12.0	17	8%	29.0	42	21%	16.2	24	12%	20.9	32	14%	19.1	30	15%	768	10%
Hispanic/Latino Males	42.5	14	7%	38.5	13	6%	28.7	10	5%	32.7	12	5%	30.7	12	6%	251	3%
Hispanic/Latina Females	*	2	1%	*	-	-	*	-	-	*	1	<1%	*	2	1%	42	1%
White Males	21.2	84	39%	12.8	51	25%	14.6	58	30%	18.2	73	33%	12.2	49	24%	3,757	50%
White Females	*	3	1%	2.2	9	4%	2.2	9	5%	2.9	12	5%	3.6	15	7%	319	4%
Multi-Race Males	31.0	5	2%	*	4	2%	39.5	7	4%	32.5	6	3%	*	3	1%	183	2%
Multi-Race Females	*	1	<1%	*	1	<1%	*	1	1%	*	1	<1%	*	3	1%	34	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	22	<1%
Total	17.5	215		16.1	202		15.5	196		17.3	224		15.5	203		7,491	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by transmission category, Franklin County

Transmission Category ^a	Diagnosis of HIV Infection by Year											
	2014		2015		2016		2017		2018		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	149	78%	108	74%	110	68%	110	63%	102	68%	4,548	72%
Injection drug use (IDU)	1	1%	2	1%	5	3%	9	5%	6	4%	216	3%
Male-to-male sexual contact and IDU	9	5%	4	3%	6	4%	8	5%	8	5%	238	4%
Heterosexual contact	5	3%	2	1%	6	4%	3	2%	4	3%	260	4%
Other/unknown	27	14%	30	21%	34	21%	46	26%	30	20%	1,022	16%
Subtotal	191	100%	146	100%	161	100%	176	100%	150	100%	6,284	100%
Female adult or adolescent												
Injection drug use	-	-	8	15%	5	15%	9	19%	11	22%	142	12%
Heterosexual contact	22	96%	43	80%	29	85%	38	79%	36	73%	884	76%
Other/unknown	1	4%	3	6%	-	-	1	2%	2	4%	130	11%
Subtotal	23	100%	54	100%	34	100%	48	100%	49	100%	1,156	100%
Child (<13 yrs at diagnosis)												
Perinatal	1	100%	1	50%	1	100%	-	-	4	100%	41	80%
Other/unknown	-	-	1	50%	-	-	-	-	-	-	10	20%
Subtotal	1	100%	2	100%	1	100%	-	-	4	100%	51	100%
Total	215		202		196		224		203		7,491	

Trends in reported diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by exposure category, Franklin County

Exposure Category ^b	Diagnosis of HIV Infection by Year										Cumulative Diagnoses	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	144	67%	105	52%	110	56%	108	48%	100	49%	4,388	59%
Injection drug use (IDU) only	-	-	4	2%	5	3%	10	4%	8	4%	223	3%
Heterosexual contact only	27	13%	45	22%	35	18%	41	18%	40	20%	1,144	15%
Male-to-male sexual contact & IDU	6	3%	4	2%	5	3%	6	3%	8	4%	202	3%
IDU & Heterosexual contact	1	<1%	6	3%	5	3%	8	4%	9	4%	135	2%
Male-to-male sexual contact & Heterosexual contact	5	2%	3	1%	-	-	2	1%	2	1%	160	2%
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	-	-	1	1%	2	1%	-	-	36	<1%
Perinatal exposure	1	<1%	1	<1%	1	1%	-	-	4	2%	43	1%
Other/unknown	28	13%	34	17%	34	17%	47	21%	32	16%	1,160	15%
Total	215		202		196		224		203		7,491	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Persons Living with Diagnosed HIV Infection Reported in Franklin County

As of June 30, 2019

Reported persons living with diagnosed HIV infection in 2018 by current disease status and selected characteristics, Franklin County

Characteristic	Living with diagnosed HIV infection			Current Disease Status			
	in 2018			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	654.6	4,190	80%	2,287	80%	1,903	80%
Females	157.7	1,057	20%	586	20%	471	20%
Age at end of year							
<13	5.3	12	<1%	7	<1%	5	<1%
13-14	*	4	<1%	3	<1%	1	<1%
15-19	28.8	24	<1%	18	1%	6	<1%
20-24	151.4	138	3%	117	4%	21	1%
25-29	387.7	492	9%	373	13%	119	5%
30-34	528.4	598	11%	393	14%	205	9%
35-39	586.5	559	11%	328	11%	231	10%
40-44	649.8	520	10%	283	10%	237	10%
45-49	821.1	655	12%	325	11%	330	14%
50-54	1,082.7	819	16%	379	13%	440	19%
55-64	746.8	1,112	21%	500	17%	612	26%
65+	199.5	314	6%	147	5%	167	7%
Race/Ethnicity^b							
American Indian/Alaska Native	202.0	5	<1%	4	<1%	1	<1%
Asian/Pacific Islander	56.6	42	1%	29	1%	13	1%
Black/African-American	771.0	2,314	44%	1,260	44%	1,054	44%
Hispanic/Latinx	421.0	315	6%	168	6%	147	6%
White	283.5	2,324	44%	1,293	45%	1,031	43%
Multi-Race	585.5	227	4%	99	3%	128	5%
Unknown	*	20	<1%	20	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	*	4	<1%	4	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	87.6	32	1%	21	1%	11	<1%
Asian/Pacific Islander Females	26.5	10	<1%	8	<1%	2	<1%
Black/African-American Males	1,109.3	1,586	30%	854	30%	732	31%
Black/African-American Females	463.3	728	14%	406	14%	322	14%
Hispanic/Latino Males	663.4	259	5%	141	5%	118	5%
Hispanic/Latina Females	156.5	56	1%	27	1%	29	1%
White Males	525.3	2,108	40%	1,174	41%	934	39%
White Females	51.6	216	4%	119	4%	97	4%
Multi-Race Males	965.5	184	4%	76	3%	108	5%
Multi-Race Females	218.1	43	1%	23	1%	20	1%
Unknown	*	20	<1%	20	1%	-	-
Total	400.4	5,247		2,873		2,374	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by current disease status and transmission category, Franklin County

Transmission Category ^a	Living with diagnosed HIV infection in 2018		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3,007	72%	1,677	74%	1,330	70%
Injection drug use (IDU)	101	2%	40	2%	61	3%
Male-to-male sexual contact and IDU	169	4%	84	4%	85	4%
Heterosexual contact	227	5%	76	3%	151	8%
Other/unknown	667	16%	397	17%	270	14%
Subtotal	4,171	100%	2,274	100%	1,897	100%
Female adult or adolescent						
Injection drug use	97	9%	41	7%	56	12%
Heterosexual contact	838	81%	458	80%	380	83%
Other/unknown	101	10%	77	13%	24	5%
Subtotal	1,036	100%	576	100%	460	100%
Child (<13 yrs at diagnosis)						
Perinatal	37	93%	21	91%	16	94%
Other/unknown	3	8%	2	9%	1	6%
Subtotal	40	100%	23	100%	17	100%
Total	5,247		2,873		2,374	

Reported persons living with diagnosed HIV infection in 2018 by current disease status and exposure category, Franklin County

Exposure Category ^b	Living with diagnosed HIV infection in 2018		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,877	55%	1,618	56%	1,259	53%
Injection drug use (IDU) only	98	2%	38	1%	60	3%
Heterosexual contact only	1,066	20%	535	19%	531	22%
Male-to-male sexual contact & IDU	140	3%	68	2%	72	3%
IDU & Heterosexual contact	100	2%	43	1%	57	2%
Male-to-male sexual contact & Heterosexual contact	130	2%	59	2%	71	3%
Male-to-male sexual contact & IDU & Heterosexual contact	29	1%	16	1%	13	1%
Perinatal exposure	39	1%	22	1%	17	1%
Other/unknown	768	15%	474	16%	294	12%
Total	5,247		2,873		2,374	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and age, Franklin County

Living with diagnosed HIV infection in 2018

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	2	5%	*	7	<1%	*	2	1%	*	-	-	*	1	<1%	-	-
13-14	*	-	-	*	1	2%	*	2	<1%	*	1	<1%	*	-	-	*	-	-	-	-
15-19	*	-	-	*	2	5%	85.5	18	1%	*	-	-	*	4	<1%	*	-	-	-	-
20-24	*	1	20%	*	2	5%	421.9	91	4%	229.2	14	4%	47.0	25	1%	133.4	5	2%	-	-
25-29	*	-	-	77.4	7	17%	1,019.0	282	12%	403.3	27	9%	181.1	144	6%	860.7	32	14%	-	-
30-34	*	-	-	*	3	7%	1,379.5	320	14%	855.9	55	17%	275.0	197	8%	915.6	23	10%	-	-
35-39	*	-	-	*	3	7%	1,252.0	263	11%	633.3	38	12%	374.6	222	10%	1,705.8	32	14%	1	5%
40-44	*	1	20%	*	4	10%	1,376.5	252	11%	800.9	41	13%	398.8	198	9%	1,524.2	23	10%	1	5%
45-49	*	2	40%	198.9	9	21%	1,431.1	263	11%	1,025.1	40	13%	609.4	314	14%	2,031.3	26	11%	1	5%
50-54	*	-	-	168.3	6	14%	1,694.9	280	12%	1,230.8	36	11%	900.5	464	20%	3,184.7	30	13%	3	15%
55-64	*	1	20%	*	2	5%	1,340.3	413	18%	1,332.2	48	15%	557.8	595	26%	2,788.6	43	19%	10	50%
65+	*	-	-	*	1	2%	454.8	123	5%	540.3	13	4%	132.5	161	7%	1,091.9	12	5%	4	20%
Total	202.0	5		56.6	42		771.0	2,314		421.0	315		283.5	2,324		585.5	227		20	

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and transmission category, Franklin County

Living with diagnosed HIV infection in 2018

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	4	100%	17	57%	970	62%	173	68%	1,711	81%	131	71%	1	6%
Injection drug use (IDU)	-	-	-	-	57	4%	5	2%	34	2%	5	3%	-	-
Male-to-male sexual contact and IDU	-	-	1	3%	45	3%	11	4%	96	5%	16	9%	-	-
Heterosexual contact	-	-	1	3%	152	10%	23	9%	42	2%	9	5%	-	-
Other/unknown	-	-	11	37%	350	22%	43	17%	224	11%	23	13%	16	94%
Subtotal	4	100%	30	100%	1,574	100%	255	100%	2,107	100%	184	100%	17	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	50	7%	3	5%	40	19%	4	10%	-	-
Heterosexual contact	1	100%	3	38%	599	84%	51	93%	152	71%	32	76%	-	-
Other/unknown	-	-	5	63%	64	9%	1	2%	22	10%	6	14%	3	100%
Subtotal	1	100%	8	100%	713	100%	55	100%	214	100%	42	100%	3	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	4	100%	24	89%	5	100%	3	100%	1	100%	-	-
Other/unknown	-	-	-	-	3	11%	-	-	-	-	-	-	-	-
Subtotal	-	-	4	100%	27	100%	5	100%	3	100%	1	100%	-	-
Total	5		42		2,314		315		2,324		227		20	

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and exposure category, Franklin County

Living with diagnosed HIV infection in 2018

Exposure Category ^a	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	4	80%	16	38%	894	39%	164	52%	1,675	72%	123	54%	1	5%
Injection drug use (IDU) only	-	-	-	-	55	2%	7	2%	33	1%	3	1%	-	-
Heterosexual contact only	1	20%	4	10%	751	32%	74	23%	195	8%	41	18%	-	-
Male-to-male sexual contact & IDU	-	-	1	2%	37	2%	11	3%	78	3%	13	6%	-	-
IDU & Heterosexual contact	-	-	-	-	52	2%	1	0%	41	2%	6	3%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	2%	76	3%	9	3%	36	2%	8	4%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	8	0%	-	-	18	1%	3	1%	-	-
Perinatal exposure	-	-	4	10%	25	1%	5	2%	3	0%	2	1%	-	-
Other/unknown	-	-	16	38%	416	18%	44	14%	245	11%	28	12%	19	95%
Total	5		42		2,314		315		2,324		227		20	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, Franklin County, 2014-2018

Characteristic	Living with Diagnosed HIV Infection														
	2014			2015			2016			2017			2018		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	638.8	3,833	82%	634.3	3,870	81%	648.8	4,004	81%	655.9	4,134	81%	654.6	4,190	80%
Females	133.8	845	18%	141.8	910	19%	147.2	953	19%	151.0	999	19%	157.7	1,057	20%
Age at end of year															
<13	6.0	13	<1%	6.4	14	<1%	5.4	12	<1%	4.9	11	<1%	5.3	12	<1%
13-14	*	3	<1%	*	4	<1%	*	4	<1%	*	4	<1%	*	4	<1%
15-19	22.9	18	<1%	22.6	18	<1%	29.8	24	<1%	31.2	26	1%	28.8	24	<1%
20-24	256.8	238	5%	217.3	195	4%	182.3	163	3%	162.8	148	3%	151.4	138	3%
25-29	384.1	455	10%	385.2	474	10%	378.7	475	10%	408.2	517	10%	387.7	492	9%
30-34	448.9	447	10%	450.7	458	10%	489.8	510	10%	495.6	536	10%	528.4	598	11%
35-39	564.6	475	10%	581.7	508	11%	603.8	543	11%	610.1	566	11%	586.5	559	11%
40-44	745.3	598	13%	685.8	548	11%	675.9	528	11%	668.9	527	10%	649.8	520	10%
45-49	1,015.2	777	17%	968.3	754	16%	902.0	715	14%	825.3	663	13%	821.1	655	12%
50-54	947.4	767	16%	990.9	795	17%	1,054.1	819	17%	1,078.1	825	16%	1,082.7	819	16%
55-64	524.8	731	16%	575.4	824	17%	645.3	933	19%	705.5	1,039	20%	746.8	1,112	21%
65+	116.6	156	3%	135.7	188	4%	161.2	231	5%	179.0	271	5%	199.5	314	6%
Race/Ethnicity^b															
American Indian/Alaska Native	*	4	<1%	*	4	<1%	*	4	<1%	202.0	5	<1%	202.0	5	<1%
Asian/Pacific Islander	44.6	25	1%	53.6	32	1%	59.3	38	1%	59.2	41	1%	56.6	42	1%
Black/African-American	711.9	1,902	41%	722.5	1,985	42%	747.9	2,109	43%	762.3	2,227	43%	771.0	2,314	44%
Hispanic/Latinx	397.4	250	5%	390.7	254	5%	407.6	274	6%	416.2	295	6%	421.0	315	6%
White	279.6	2,263	48%	277.3	2,260	47%	281.1	2,284	46%	282.8	2,318	45%	283.5	2,324	44%
Multi-Race	640.0	214	5%	648.5	225	5%	628.5	228	5%	603.9	227	4%	585.5	227	4%
Unknown	*	20	<1%	*	20	<1%	*	20	<1%	*	20	<1%	*	20	<1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	3	<1%	*	3	<1%	*	3	<1%	*	4	<1%	*	4	<1%
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	76.0	21	<1%	81.6	24	1%	91.2	29	1%	94.0	32	1%	87.6	32	1%
Asian/Pacific Islander Females	*	4	<1%	26.4	8	<1%	27.9	9	<1%	25.5	9	<1%	26.5	10	<1%
Black/African-American Males	1,061.7	1,337	29%	1,055.7	1,369	29%	1,089.6	1,458	29%	1,111.2	1,541	30%	1,109.3	1,586	30%
Black/African-American Females	400.1	565	12%	424.7	616	13%	439.4	651	13%	447.0	686	13%	463.3	728	14%
Hispanic/Latino Males	606.9	200	4%	601.0	203	4%	635.3	221	4%	653.1	240	5%	663.4	259	5%
Hispanic/Latina Females	166.9	50	1%	163.2	51	1%	163.4	53	1%	161.1	55	1%	156.5	56	1%
White Males	523.0	2,072	44%	517.4	2,066	43%	524.1	2,085	42%	526.4	2,112	41%	525.3	2,108	40%
White Females	46.2	191	4%	46.7	194	4%	48.0	199	4%	49.2	206	4%	51.6	216	4%
Multi-Race Males	1,134.7	183	4%	1,122.8	188	4%	1,078.4	191	4%	1,019.6	188	4%	965.5	184	4%
Multi-Race Females	179.1	31	1%	206.1	37	1%	199.3	37	1%	203.6	39	1%	218.1	43	1%
Unknown	*	20	<1%	*	20	<1%	*	20	<1%	*	20	<1%	*	20	<1%
Total	379.9	4,678		381.9	4,780		392.0	4,957		397.3	5,133		400.4	5,247	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-Race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, Franklin County, 2014-2018

Living with Diagnosed HIV Infection										
Transmission Category ^a	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	2,799	73%	2,814	73%	2,903	73%	2,981	72%	3,007	72%
Injection drug use (IDU)	82	2%	78	2%	86	2%	93	2%	101	2%
Male-to-male sexual contact and IDU	160	4%	162	4%	165	4%	172	4%	169	4%
Heterosexual contact	212	6%	209	5%	214	5%	215	5%	227	5%
Other/unknown	565	15%	591	15%	617	15%	654	16%	667	16%
Subtotal	3,818	100%	3,854	100%	3,985	100%	4,115	100%	4,171	100%
Female adult or adolescent										
Injection drug use	73	9%	82	9%	86	9%	92	9%	97	9%
Heterosexual contact	659	80%	712	80%	752	81%	791	81%	838	81%
Other/unknown	92	11%	94	11%	94	10%	95	10%	101	10%
Subtotal	824	100%	888	100%	932	100%	978	100%	1,036	100%
Child (<13 yrs at diagnosis)										
Perinatal	32	89%	34	89%	35	88%	36	90%	37	93%
Other/unknown	4	11%	4	11%	5	13%	4	10%	3	8%
Subtotal	36	100%	38	100%	40	100%	40	100%	40	100%
Total	4,678		4,780		4,957		5,133		5,247	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, Franklin County, 2014-2018

Living with Diagnosed HIV Infection										
Exposure Category ^b	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,649	57%	2,670	56%	2,762	56%	2,839	55%	2,877	55%
Injection drug use (IDU) only	73	2%	77	2%	82	2%	85	2%	98	2%
Heterosexual contact only	872	19%	922	19%	967	20%	1,007	20%	1,066	20%
Male-to-male sexual contact & IDU	127	3%	128	3%	130	3%	137	3%	140	3%
IDU & Heterosexual contact	82	2%	83	2%	90	2%	100	2%	100	2%
Male-to-male sexual contact & Heterosexual contact	150	3%	144	3%	141	3%	142	3%	130	2%
Male-to-male sexual contact & IDU & Heterosexual contact	33	1%	34	1%	35	1%	35	1%	29	1%
Perinatal exposure	34	1%	36	1%	37	1%	38	1%	39	1%
Other/unknown	658	14%	686	14%	713	14%	750	15%	768	15%
Total	4,678		4,780		4,957		5,133		5,247	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.