

New Diagnoses of HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A- Columbus

Counties: Delaware, Fairfield, Franklin, Licking, Madison, Morrow, Pickaway, Union

Reported new diagnoses of HIV infection in 2018 by disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	2018 diagnosis of HIV infection			Disease Status					
	Rate ^a	No.	%	HIV (not AIDS)		HIV & later AIDS		AIDS	
				No.	%	No.	%	No.	%
Sex at birth									
Males	17.1	172	76%	131	77%	29	69%	12	86%
Females	5.2	54	24%	39	23%	13	31%	2	14%
Age at diagnosis (yr)									
<13	*	4	2%	3	2%	-	-	1	7%
13-14	*	-	-	-	-	-	-	-	-
15-19	9.0	12	5%	10	6%	2	5%	-	-
20-24	23.2	31	14%	27	16%	2	5%	2	14%
25-29	32.2	54	24%	47	28%	6	14%	1	7%
30-34	23.5	37	16%	28	16%	7	17%	2	14%
35-39	14.5	21	9%	12	7%	6	14%	3	21%
40-44	16.3	21	9%	14	8%	5	12%	2	14%
45-49	14.2	19	8%	14	8%	5	12%	-	-
50-54	8.7	11	5%	6	4%	5	12%	-	-
55-64	4.9	12	5%	6	4%	4	10%	2	14%
65+	*	4	2%	3	2%	-	-	1	7%
Race/Ethnicity^b									
American Indian/Alaska Native	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander	*	2	1%	2	1%	-	-	-	-
Black/African-American	37.1	124	55%	95	56%	23	55%	6	43%
Hispanic/Latinx	15.4	14	6%	10	6%	2	5%	2	14%
White	5.5	80	35%	59	35%	16	38%	5	36%
Multi-Race	11.4	6	3%	4	2%	1	2%	1	7%
Race/Ethnicity^b and Sex at birth									
American Indian/Alaska Native Males	*	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	*	-	-	-	-	-	-	-	-
Asian/Pacific Islander Males	*	1	<1%	1	1%	-	-	-	-
Asian/Pacific Islander Females	*	1	<1%	1	1%	-	-	-	-
Black/African-American Males	58.0	94	42%	72	42%	16	38%	6	43%
Black/African-American Females	17.5	30	13%	23	14%	7	17%	-	-
Hispanic/Latino Males	25.4	12	5%	8	5%	2	5%	2	14%
Hispanic/Latina Females	*	2	1%	2	1%	-	-	-	-
White Males	8.6	62	27%	48	28%	11	26%	3	21%
White Females	2.4	18	8%	11	6%	5	12%	2	14%
Multi-Race Males	*	3	1%	2	1%	-	-	1	7%
Multi-Race Females	*	3	1%	2	1%	1	2%	-	-
Total	11.1	226		170		42		14	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^aThe rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.

^bHispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent								
Male-to-male sexual contact	110	65%	88	68%	16	55%	6	55%
Injection drug use (IDU)	7	4%	5	4%	1	3%	1	9%
Male-to-male sexual contact and IDU	9	5%	9	7%	-	-	-	-
Heterosexual contact	7	4%	4	3%	2	7%	1	9%
Other/unknown	37	22%	24	18%	10	34%	3	27%
Subtotal	170	100%	130	100%	29	100%	11	100%
Female adult or adolescent								
Injection drug use	12	23%	6	16%	5	38%	1	50%
Heterosexual contact	37	71%	29	78%	7	54%	1	50%
Other/unknown	3	6%	2	5%	1	8%	-	-
Subtotal	52	100%	37	100%	13	100%	2	100%
Child (<13 yrs at diagnosis)								
Perinatal	4	100%	3	100%	-	-	1	100%
Other/unknown	-	-	-	-	-	-	-	-
Subtotal	4	100%	3	100%	-	-	1	100%
Total	226		170		42		14	

Reported new diagnoses of HIV infection in 2018 by disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	2018 diagnosis of HIV infection		Disease Status					
			HIV (not AIDS)		HIV & later AIDS		AIDS	
	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	108	48%	86	51%	16	38%	6	43%
Injection drug use (IDU) only	8	4%	5	3%	3	7%	-	-
Heterosexual contact only	44	19%	33	19%	9	21%	2	14%
Male-to-male sexual contact & IDU	9	4%	9	5%	-	-	-	-
IDU & Heterosexual contact	11	5%	6	4%	3	7%	2	14%
Male-to-male sexual contact & Heterosexual contact	2	1%	2	1%	-	-	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-
Perinatal exposure	4	2%	3	2%	-	-	1	7%
Other/unknown	40	18%	26	15%	11	26%	3	21%
Total	226		170		42		14	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and age at diagnosis, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

2018 diagnosis of HIV infection																		
Age at diagnosis (yr)	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race		
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%
<13	*	-	-	*	1	50%	*	3	2%	*	-	-	*	-	-	*	-	-
13-14	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-
15-19	*	-	-	*	1	50%	33.4	8	6%	*	-	-	*	3	4%	*	-	-
20-24	*	-	-	*	-	-	77.6	19	15%	*	4	29%	7.9	7	9%	*	1	17%
25-29	*	-	-	*	-	-	115.4	35	28%	*	2	14%	14.9	17	21%	*	-	-
30-34	*	-	-	*	-	-	58.3	15	12%	65.5	5	36%	13.8	15	19%	*	2	33%
35-39	*	-	-	*	-	-	46.4	11	9%	*	1	7%	7.9	8	10%	*	1	17%
40-44	*	-	-	*	-	-	33.3	7	6%	*	1	7%	14.2	13	16%	*	-	-
45-49	*	-	-	*	-	-	47.3	10	8%	*	1	7%	7.1	7	9%	*	1	17%
50-54	*	-	-	*	-	-	42.5	8	6%	*	-	-	*	3	4%	*	-	-
55-64	*	-	-	*	-	-	17.6	6	5%	*	-	-	3.1	6	8%	*	-	-
65+	*	-	-	*	-	-	*	2	2%	*	-	-	*	1	1%	*	1	17%
Total	*	-	-	*	2	-	37.1	124	-	15.4	14	-	5.5	80	-	11.4	6	-

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^c	2018 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	-	-	-	-	56	61%	10	83%	42	68%	2	67%
Injection drug use (IDU)	-	-	-	-	1	1%	-	-	6	10%	-	-
Male-to-male sexual contact and IDU	-	-	-	-	3	3%	-	-	6	10%	-	-
Heterosexual contact	-	-	-	-	4	4%	1	8%	2	3%	-	-
Other/unknown	-	-	1	100%	28	30%	1	8%	6	10%	1	33%
Subtotal	-	-	1	100%	92	100%	12	100%	62	100%	3	100%
Female adult or adolescent												
Injection drug use	-	-	-	-	1	3%	-	-	11	61%	-	-
Heterosexual contact	-	-	-	-	27	93%	2	100%	6	33%	2	67%
Other/unknown	-	-	-	-	1	3%	-	-	1	6%	1	33%
Subtotal	-	-	-	-	29	100%	2	100%	18	100%	3	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	1	100%	3	100%	-	-	-	-	-	-
Other/unknown	-	-	-	-	-	-	-	-	-	-	-	-
Subtotal	-	-	1	100%	3	100%	-	-	-	-	-	-
Total	-		2		124		14		80		6	

Reported new diagnoses of HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^d	2018 diagnosis of HIV infection											
	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/Latinx ^a		White		Multi-Race	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	-	-	-	-	56	45%	9	64%	41	51%	2	33%
Injection drug use (IDU) only	-	-	-	-	2	2%	-	-	6	8%	-	-
Heterosexual contact only	-	-	-	-	31	25%	3	21%	8	10%	2	33%
Male-to-male sexual contact & IDU	-	-	-	-	3	2%	-	-	6	8%	-	-
IDU & Heterosexual contact	-	-	-	-	-	-	-	-	11	14%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	-	-	-	-	1	7%	1	1%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	-	-	1	50%	3	2%	-	-	-	-	-	-
Other/unknown	-	-	1	50%	29	23%	1	7%	7	9%	2	33%
Total	-	-	2	-	124	-	14	-	80	-	6	-

Notes:
Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis.
Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.
^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.
^b The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using 2018 U.S. Census estimates.
^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.
^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.
Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Diagnosis of HIV Infection by Year																	
Characteristic	2014			2015			2016			2017			2018			Cumulative Diagnoses	
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	No.	%
Sex at birth																	
Males	21.2	201	87%	17.5	168	75%	18.2	177	81%	19.6	194	79%	17.1	172	76%	6,995	84%
Females	3.1	30	13%	5.5	55	25%	4.2	42	19%	5.0	51	21%	5.2	54	24%	1,299	16%
Age at diagnosis (yr)																	
<13	*	1	<1%	*	2	1%	*	1	<1%	*	-	-	*	4	2%	59	1%
13-14	*	-	-	*	-	-	*	1	<1%	*	-	-	*	-	-	10	<1%
15-19	5.5	7	3%	11.6	15	7%	10.8	14	6%	8.3	11	4%	9.0	12	5%	293	4%
20-24	36.7	49	21%	29.7	39	17%	29.0	38	17%	27.1	36	15%	23.2	31	14%	1,060	13%
25-29	31.5	49	21%	26.2	42	19%	24.4	40	18%	34.3	57	23%	32.2	54	24%	1,467	18%
30-34	25.3	36	16%	21.4	31	14%	16.9	25	11%	23.1	35	14%	23.5	37	16%	1,583	19%
35-39	20.7	27	12%	20.9	28	13%	19.6	27	12%	19.0	27	11%	14.5	21	9%	1,377	17%
40-44	13.6	18	8%	14.6	19	9%	16.5	21	10%	14.9	19	8%	16.3	21	9%	1,009	12%
45-49	18.8	24	10%	16.9	22	10%	11.3	15	7%	11.9	16	7%	14.2	19	8%	670	8%
50-54	6.0	8	3%	6.7	9	4%	15.4	20	9%	14.1	18	7%	8.7	11	5%	396	5%
55-64	4.4	10	4%	3.8	9	4%	6.7	16	7%	9.1	22	9%	4.9	12	5%	301	4%
65+	*	2	1%	2.9	7	3%	*	1	<1%	*	4	2%	*	4	2%	69	1%
Race/Ethnicity ^b																	
American Indian/Alaska Native	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-	7	<1%
Asian/Pacific Islander	*	4	2%	9.1	7	3%	8.4	7	3%	*	3	1%	*	2	1%	53	1%
Black/African-American	35.0	104	45%	39.3	120	54%	35.8	112	51%	36.9	120	49%	37.1	124	55%	2,960	36%
Hispanic/Latinx	23.6	18	8%	16.5	13	6%	14.7	12	5%	15.1	13	5%	15.4	14	6%	312	4%
White	6.8	98	42%	5.3	76	34%	5.5	79	36%	6.8	99	40%	5.5	80	35%	4,699	57%
Multi-Race	15.5	7	3%	14.9	7	3%	18.4	9	4%	17.7	9	4%	11.4	6	3%	239	3%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	24	<1%
Race/Ethnicity ^b and Sex at birth																	
American Indian/Alaska Native Males	*	-	-	*	-	-	*	-	-	*	1	<1%	*	-	-	5	<1%
American Indian/Alaska Native Females	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	2	<1%
Asian/Pacific Islander Males	*	3	1%	*	4	2%	14.6	6	3%	*	1	<1%	*	1	<1%	38	<1%
Asian/Pacific Islander Females	*	1	<1%	*	3	1%	*	1	<1%	*	2	1%	*	1	<1%	15	<1%
Black/African-American Males	60.2	86	37%	53.1	78	35%	56.8	86	39%	56.0	88	36%	58.0	94	42%	2,162	26%
Black/African-American Females	11.7	18	8%	26.5	42	19%	16.1	26	12%	19.1	32	13%	17.5	30	13%	798	10%
Hispanic/Latino Males	37.7	15	6%	31.8	13	6%	26.1	11	5%	27.0	12	5%	25.4	12	5%	265	3%
Hispanic/Latina Females	*	3	1%	*	-	-	*	1	<1%	*	1	<1%	*	2	1%	47	1%
White Males	12.9	91	39%	9.4	67	30%	9.3	66	30%	11.8	85	35%	8.6	62	27%	4,304	52%
White Females	1.0	7	3%	1.2	9	4%	1.8	13	6%	1.9	14	6%	2.4	18	8%	395	5%
Multi-Race Males	27.2	6	3%	26.2	6	3%	33.3	8	4%	28.0	7	3%	*	3	1%	201	2%
Multi-Race Females	*	1	<1%	*	1	<1%	*	1	<1%	*	2	1%	*	3	1%	38	<1%
Unknown	*	-	-	*	-	-	*	-	-	*	-	-	*	-	-	24	<1%
Total	12.0	231		11.4	223		11.1	219		12.2	245		11.1	226		8,294	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons with a reported diagnosis of HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported new diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Diagnosis of HIV Infection by Year												Cumulative Diagnoses	
	2014		2015		2016		2017		2018					
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%		
Male adult or adolescent														
Male-to-male sexual contact	154	77%	120	72%	117	66%	120	62%	110	65%	4,978	71%		
Injection drug use (IDU)	1	<1%	4	2%	7	4%	9	5%	7	4%	259	4%		
Male-to-male sexual contact and IDU	10	5%	5	3%	8	5%	9	5%	9	5%	275	4%		
Heterosexual contact	6	3%	2	1%	6	3%	5	3%	7	4%	297	4%		
Other/unknown	30	15%	36	22%	38	22%	51	26%	37	22%	1,157	17%		
Subtotal	201	100%	167	100%	176	100%	194	100%	170	100%	6,966	100%		
Female adult or adolescent														
Injection drug use	1	3%	8	15%	6	14%	9	18%	12	23%	158	12%		
Heterosexual contact	27	93%	43	80%	34	81%	41	80%	37	71%	961	76%		
Other/unknown	1	3%	3	6%	2	5%	1	2%	3	6%	147	12%		
Subtotal	29	100%	54	100%	42	100%	51	100%	52	100%	1,266	100%		
Child (<13 yrs at diagnosis)														
Perinatal	1	100%	1	50%	1	100%	-	-	4	100%	51	82%		
Other/unknown	-	-	1	50%	-	-	-	-	-	-	11	18%		
Subtotal	1	100%	2	100%	1	100%	-	-	4	100%	62	100%		
Total	231		223		219		245		226		8,294			

Trends in reported diagnoses of HIV infection by year of diagnosis (2014-2018) and cumulative diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^p	Diagnosis of HIV Infection by Year											
	2014		2015		2016		2017		2018		Cumulative Diagnoses	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	149	65%	116	52%	117	53%	118	48%	108	48%	4,797	58%
Injection drug use (IDU) only	1	<1%	6	3%	6	3%	10	4%	8	4%	259	3%
Heterosexual contact only	33	14%	45	20%	40	18%	46	19%	44	19%	1,258	15%
Male-to-male sexual contact & IDU	7	3%	4	2%	7	3%	7	3%	9	4%	235	3%
IDU & Heterosexual contact	1	<1%	6	3%	7	3%	8	3%	11	5%	158	2%
Male-to-male sexual contact & Heterosexual contact	5	2%	4	2%	-	-	2	1%	2	1%	181	2%
Male-to-male sexual contact & IDU & Heterosexual contact	3	1%	1	<1%	1	<1%	2	1%	-	-	40	<1%
Perinatal exposure	1	<1%	1	<1%	2	1%	-	-	4	2%	54	1%
Other/unknown	31	13%	40	18%	39	18%	52	21%	40	18%	1,312	16%
Total	231		223		219		245		226		8,294	

Notes:

Reported new diagnoses of HIV infection include persons with a diagnosis of HIV (not AIDS), a diagnosis of HIV and an AIDS diagnosis within 12 months (HIV & later AIDS), and concurrent diagnoses of HIV and AIDS (AIDS) who were residents of Ohio at time of initial diagnosis. Diagnoses of HIV infection by year (2014-2018) represent all reported cases diagnosed in each year; cumulative diagnoses represent all reported HIV and/or AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Sex at birth						
Males	73	75%	30	77%	43	74%
Females	24	25%	9	23%	15	26%
Age at AIDS diagnosis (yr)						
<13	1	1%	-	-	1	2%
13-14	-	-	-	-	-	-
15-19	2	2%	1	3%	1	2%
20-24	8	8%	2	5%	6	10%
25-29	9	9%	2	5%	7	12%
30-34	19	20%	9	23%	10	17%
35-39	14	14%	6	15%	8	14%
40-44	12	12%	5	13%	7	12%
45-49	10	10%	4	10%	6	10%
50-54	10	10%	5	13%	5	9%
55-64	11	11%	5	13%	6	10%
65+	1	1%	-	-	1	2%
Race/Ethnicity^b						
American Indian/Alaska Native	-	-	-	-	-	-
Asian/Pacific Islander	-	-	-	-	-	-
Black/African-American	50	52%	22	56%	28	48%
Hispanic/Latinx	6	6%	1	3%	5	9%
White	39	40%	16	41%	23	40%
Multi-Race	2	2%	-	-	2	3%
Race/Ethnicity^b and Sex at birth						
American Indian/Alaska Native Males	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-
Asian/Pacific Islander Males	-	-	-	-	-	-
Asian/Pacific Islander Females	-	-	-	-	-	-
Black/African-American Males	39	40%	18	46%	21	36%
Black/African-American Females	11	11%	4	10%	7	12%
Hispanic/Latino Males	6	6%	1	3%	5	9%
Hispanic/Latina Females	-	-	-	-	-	-
White Males	27	28%	11	28%	16	28%
White Females	12	12%	5	13%	7	12%
Multi-Race Males	1	1%	-	-	1	2%
Multi-Race Females	1	1%	-	-	1	2%
Total	97		39		58	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^b	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12 Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	37	51%	15	50%	22	52%
Injection drug use (IDU)	6	8%	3	10%	3	7%
Male-to-male sexual contact and IDU	3	4%	3	10%	-	-
Heterosexual contact	5	7%	2	7%	3	7%
Other/unknown	21	29%	7	23%	14	33%
Subtotal	72	100%	30	100%	42	100%
Female adult or adolescent						
Injection drug use	12	50%	6	67%	6	40%
Heterosexual contact	10	42%	2	22%	8	53%
Other/unknown	2	8%	1	11%	1	7%
Subtotal	24	100%	9	100%	15	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	4%	-	-	1	7%
Other/unknown	-	-	-	-	-	-
Subtotal	1	4%	-	-	1	7%
Total	97		39		58	

Reported diagnoses of AIDS in 2018 by time to AIDS diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^c	AIDS Diagnoses					
	2018 AIDS diagnosis		≥12Months after HIV Diagnosis		<12 Months after HIV Diagnosis ^a	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	35	36%	13	33%	22	38%
Injection drug use (IDU) only	5	5%	3	8%	2	3%
Heterosexual contact only	15	15%	4	10%	11	19%
Male-to-male sexual contact & IDU	3	3%	3	8%	-	-
IDU & Heterosexual contact	13	13%	6	15%	7	12%
Male-to-male sexual contact & Heterosexual contact	2	2%	2	5%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	1%	-	-	1	2%
Other/unknown	23	24%	8	21%	15	26%
Total	97		39		58	

Notes:

Data reflects all persons with an AIDS diagnosis in 2018 regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Includes persons whose diagnoses of HIV infection and AIDS were made at the same time.

^b Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^c Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Sex at birth												
Males	96	86%	77	78%	92	82%	83	78%	73	75%	3,973	86%
Females	16	14%	22	22%	20	18%	23	22%	24	25%	640	14%
Age at AIDS diagnosis (yr)												
<13	-	-	-	-	-	-	-	-	1	1%	24	1%
13-14	-	-	-	-	-	-	-	-	-	-	4	<1%
15-19	-	-	1	1%	1	1%	1	1%	2	2%	42	1%
20-24	15	13%	7	7%	9	8%	12	11%	8	8%	273	6%
25-29	14	13%	16	16%	16	14%	7	7%	9	9%	600	13%
30-34	23	21%	14	14%	12	11%	12	11%	19	20%	946	21%
35-39	9	8%	8	8%	19	17%	15	14%	14	14%	861	19%
40-44	15	13%	12	12%	14	13%	18	17%	12	12%	757	16%
45-49	17	15%	17	17%	10	9%	10	9%	10	10%	504	11%
50-54	8	7%	16	16%	18	16%	13	12%	10	10%	323	7%
55-64	8	7%	5	5%	10	9%	16	15%	11	11%	241	5%
65+	3	3%	3	3%	3	3%	2	2%	1	1%	38	1%
Race/Ethnicity ^a												
American Indian/Alaska Native	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander	3	3%	-	-	1	1%	-	-	-	-	17	<1%
Black/African-American	50	45%	44	44%	51	46%	56	53%	50	52%	1,507	33%
Hispanic/Latinx	6	5%	8	8%	9	8%	6	6%	6	6%	163	4%
White	47	42%	44	44%	49	44%	39	37%	39	40%	2,794	61%
Multi-Race	6	5%	3	3%	2	2%	5	5%	2	2%	130	3%
Race/Ethnicity ^a and Sex at birth												
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Asian/Pacific Islander Males	3	3%	-	-	1	1%	-	-	-	-	15	<1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-	2	<1%
Black/African-American Males	39	35%	31	31%	38	34%	41	39%	39	40%	1,120	24%
Black/African-American Females	11	10%	13	13%	13	12%	15	14%	11	11%	387	8%
Hispanic/Latino Males	6	5%	7	7%	8	7%	5	5%	6	6%	140	3%
Hispanic/Latina Females	-	-	1	1%	1	1%	1	1%	-	-	23	<1%
White Males	42	38%	36	36%	43	38%	33	31%	27	28%	2,586	56%
White Females	5	4%	8	8%	6	5%	6	6%	12	12%	208	5%
Multi-Race Males	6	5%	3	3%	2	2%	4	4%	1	1%	112	2%
Multi-Race Females	-	-	-	-	-	-	1	1%	1	1%	18	<1%
Total	112		99		112		106		97		4,613	

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	65	68%	52	68%	60	65%	50	60%	37	51%	2,861	72%
Injection drug use (IDU)	-	-	2	3%	4	4%	4	5%	6	8%	174	4%
Male-to-male sexual contact and IDU	1	1%	1	1%	2	2%	7	8%	3	4%	155	4%
Heterosexual contact	9	9%	2	3%	5	5%	1	1%	5	7%	217	5%
Other/unknown	21	22%	20	26%	21	23%	21	25%	21	29%	551	14%
Subtotal	96	100%	77	100%	92	100%	83	100%	72	100%	3,958	100%
Female adult or adolescent												
Injection drug use	-	-	4	18%	3	15%	5	22%	12	50%	97	16%
Heterosexual contact	15	94%	17	77%	17	85%	15	65%	10	42%	480	77%
Other/unknown	1	6%	1	5%	-	-	3	13%	2	8%	46	7%
Subtotal	16	100%	22	100%	20	100%	23	100%	24	100%	623	100%
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	1	100%	25	78%
Other/unknown	-	-	-	-	-	-	-	-	-	-	7	22%
Subtotal	-	-	-	-	-	-	-	-	1	100%	32	100%
Total	112		99		112		106		97		4,613	

Trends in reported diagnoses of AIDS by year of diagnosis (2014-2018) and cumulative AIDS diagnoses by exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	Year of AIDS Diagnosis										Cumulative AIDS	
	2014		2015		2016		2017		2018			
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	61	54%	51	52%	58	52%	50	47%	35	36%	2,752	60%
Injection drug use (IDU) only	-	-	2	2%	4	4%	4	4%	5	5%	172	4%
Heterosexual contact only	24	21%	19	19%	22	20%	16	15%	15	15%	697	15%
Male-to-male sexual contact & IDU	1	1%	-	-	2	2%	5	5%	3	3%	133	3%
IDU & Heterosexual contact	-	-	4	4%	3	3%	5	5%	13	13%	99	2%
Male-to-male sexual contact & Heterosexual contact	4	4%	1	1%	2	2%	-	-	2	2%	109	2%
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	1	1%	-	-	2	2%	-	-	22	<1%
Perinatal exposure	-	-	-	-	-	-	-	-	1	1%	26	1%
Other/unknown	22	20%	21	21%	21	19%	24	23%	23	24%	603	13%
Total	112		99		112		106		97		4,613	

Notes:

AIDS diagnoses by year (2014-2018) represent all reported AIDS cases diagnosed in each year regardless of year of HIV infection diagnosis, and who were living in Ohio at the time of initial diagnosis; cumulative AIDS cases represent all reported AIDS cases diagnosed since the beginning of the epidemic through 2018.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Persons Living with Diagnosed HIV Infection Reported in HIV Prevention Planning Region 11/Ryan White Part A- Columbus

Counties: Delaware, Fairfield, Franklin, Licking, Madison, Morrow, Pickaway, Union

Reported persons living with diagnosed HIV infection in 2018 by current disease status and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Characteristic	Living with diagnosed HIV infection in			Current Disease Status			
	2018			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Sex at birth							
Males	465.6	4,680	80%	2,542	79%	2,138	80%
Females	114.9	1,191	20%	657	21%	534	20%
Age at end of year							
<13	4.3	15	<1%	10	<1%	5	<1%
13-14	9.5	5	<1%	3	<1%	2	<1%
15-19	21.7	29	<1%	21	1%	8	<1%
20-24	121.3	162	3%	139	4%	23	1%
25-29	317.0	531	9%	404	13%	127	5%
30-34	395.4	622	11%	411	13%	211	8%
35-39	419.2	609	10%	362	11%	247	9%
40-44	446.5	577	10%	304	10%	273	10%
45-49	565.5	755	13%	374	12%	381	14%
50-54	742.4	940	16%	439	14%	501	19%
55-64	516.5	1,265	22%	563	18%	702	26%
65+	134.6	361	6%	169	5%	192	7%
Race/Ethnicity^b							
American Indian/Alaska Native	150.7	6	<1%	5	<1%	1	<1%
Asian/Pacific Islander	49.7	49	1%	35	1%	14	1%
Black/African-American	727.0	2,427	41%	1,323	41%	1,104	41%
Hispanic/Latinx	369.1	335	6%	179	6%	156	6%
White	189.7	2,775	47%	1,520	48%	1,255	47%
Multi-Race	489.8	257	4%	115	4%	142	5%
Unknown	*	22	<1%	22	1%	-	-
Race/Ethnicity^b and Sex at birth							
American Indian/Alaska Native Males	252.5	5	<1%	5	<1%	-	-
American Indian/Alaska Native Females	*	1	<1%	-	-	1	<1%
Asian/Pacific Islander Males	74.4	36	1%	25	1%	11	<1%
Asian/Pacific Islander Females	25.9	13	<1%	10	<1%	3	<1%
Black/African-American Males	1,024.2	1,659	28%	896	28%	763	29%
Black/African-American Females	446.9	768	13%	427	13%	341	13%
Hispanic/Latino Males	573.7	271	5%	147	5%	124	5%
Hispanic/Latina Females	147.0	64	1%	32	1%	32	1%
White Males	344.8	2,482	42%	1,360	43%	1,122	42%
White Females	39.5	293	5%	160	5%	133	5%
Multi-Race Males	809.5	209	4%	91	3%	118	4%
Multi-Race Females	180.1	48	1%	24	1%	24	1%
Unknown	*	22	<1%	22	1%	-	-
Total	287.5	5,871		3,199		2,672	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by current disease status and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Transmission Category ^a	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	3,335	72%	1,845	73%	1,490	70%
Injection drug use (IDU)	111	2%	42	2%	69	3%
Male-to-male sexual contact and IDU	195	4%	95	4%	100	5%
Heterosexual contact	256	5%	87	3%	169	8%
Other/unknown	758	16%	457	18%	301	14%
Subtotal	4,655	100%	2,526	100%	2,129	100%
Female adult or adolescent						
Injection drug use	109	9%	46	7%	63	12%
Heterosexual contact	931	80%	500	78%	431	83%
Other/unknown	119	10%	93	15%	26	5%
Subtotal	1,159	100%	639	100%	520	100%
Child (<13 yrs at diagnosis)						
Perinatal	52	91%	31	91%	21	91%
Other/unknown	5	9%	3	9%	2	9%
Subtotal	57	100%	34	100%	23	100%
Total	5,871		3,199		2,672	

Reported persons living with diagnosed HIV infection in 2018 by current disease status and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Exposure Category ^b	Living with diagnosed HIV infection in		Current Disease Status			
	2018		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male-to-male sexual contact only	3,189	54%	1,777	56%	1,412	53%
Injection drug use (IDU) only	108	2%	40	1%	68	3%
Heterosexual contact only	1,188	20%	588	18%	600	22%
Male-to-male sexual contact & IDU	160	3%	75	2%	85	3%
IDU & Heterosexual contact	112	2%	48	2%	64	2%
Male-to-male sexual contact & Heterosexual contact	146	2%	68	2%	78	3%
Male-to-male sexual contact & IDU & Heterosexual contact	35	1%	20	1%	15	1%
Perinatal exposure	55	1%	33	1%	22	1%
Other/unknown	878	15%	550	17%	328	12%
Total	5,871		3,199		2,672	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and age, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2018

Age at end of year	American Indian/Alaska Native			Asian/Pacific Islander			Black/African-American			Hispanic/ Latinx ^a			White			Multi-Race			Unknown	
	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	Rate ^b	No.	%	No.	%
<13	*	-	-	*	2	4%	*	9	<1%	*	3	1%	*	-	-	*	1	<1%	-	-
13-14	*	-	-	*	1	2%	*	3	<1%	*	1	<1%	*	-	-	*	-	-	-	-
15-19	*	-	-	*	3	6%	83.6	20	1%	*	-	-	6.7	6	<1%	*	-	-	-	-
20-24	*	1	17%	*	3	6%	404.5	99	4%	233.3	17	5%	38.2	34	1%	161.8	8	3%	-	-
25-29	*	-	-	66.2	7	14%	976.1	296	12%	371.1	29	9%	142.1	162	6%	829.0	37	14%	-	-
30-34	*	-	-	*	4	8%	1,270.8	327	13%	733.8	56	17%	190.8	208	7%	872.7	27	11%	-	-
35-39	*	-	-	*	3	6%	1,148.5	272	11%	533.3	39	12%	256.3	261	9%	1,349.7	33	13%	1	5%
40-44	*	1	17%	76.5	6	12%	1,259.4	265	11%	648.5	41	12%	259.5	238	9%	1,228.5	25	10%	1	5%
45-49	*	2	33%	150.3	10	20%	1,334.3	282	12%	913.1	45	13%	390.8	386	14%	1,652.4	29	11%	1	5%
50-54	*	1	17%	123.3	6	12%	1,560.2	294	12%	971.4	36	11%	579.0	565	20%	2,548.7	34	13%	4	18%
55-64	*	1	17%	*	3	6%	1,272.5	435	18%	1,162.0	53	16%	362.9	711	26%	2,258.6	51	20%	11	50%
65+	*	-	-	*	1	2%	425.4	125	5%	464.7	15	4%	90.0	204	7%	695.2	12	5%	4	18%
Total	150.7	6		49.7	49		727.0	2,427		369.1	335		189.7	2,775		489.8	257		22	

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2018

Transmission Category ^c	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent														
Male-to-male sexual contact	5	100%	18	53%	1,008	61%	181	68%	1,974	80%	148	71%	1	6%
Injection drug use (IDU)	-	-	-	-	59	4%	6	2%	41	2%	5	2%	-	-
Male-to-male sexual contact and IDU	-	-	2	6%	48	3%	11	4%	117	5%	17	8%	-	-
Heterosexual contact	-	-	2	6%	160	10%	23	9%	60	2%	11	5%	-	-
Other/unknown	-	-	12	35%	371	23%	46	17%	286	12%	26	13%	17	94%
Subtotal	5	100%	34	100%	1,646	100%	267	100%	2,478	100%	207	100%	18	100%
Female adult or adolescent														
Injection drug use	-	-	-	-	51	7%	4	7%	49	17%	5	11%	-	-
Heterosexual contact only	1	100%	5	45%	633	85%	55	92%	202	70%	35	74%	-	-
Other/unknown	-	-	6	55%	64	9%	1	2%	37	13%	7	15%	4	100%
Subtotal	1	100%	11	100%	748	100%	60	100%	288	100%	47	100%	4	100%
Child (<13 yrs at diagnosis)														
Perinatal	-	-	4	100%	30	91%	8	100%	7	78%	3	100%	-	-
Other/unknown	-	-	-	-	3	9%	-	-	2	22%	-	-	-	-
Subtotal	-	-	4	100%	33	100%	8	100%	9	100%	3	100%	-	-
Total	6		49		2,427		335		2,775		257		22	

Reported persons living with diagnosed HIV infection in 2018 by race/ethnicity and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus

Living with diagnosed HIV infection in 2018

Exposure Category ^d	American Indian/Alaska Native		Asian/Pacific Islander		Black/African-American		Hispanic/ Latinx ^a		White		Multi-Race		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	5	83%	17	35%	928	38%	170	51%	1,929	70%	139	54%	1	5%
Injection drug use (IDU) only	-	-	-	-	57	2%	9	3%	38	1%	4	2%	-	-
Heterosexual contact only	1	17%	7	14%	793	33%	78	23%	263	9%	46	18%	-	-
Male-to-male sexual contact & IDU	-	-	2	4%	39	2%	11	3%	94	3%	14	5%	-	-
IDU & Heterosexual contact	-	-	-	-	53	2%	1	0%	52	2%	6	2%	-	-
Male-to-male sexual contact & Heterosexual contact	-	-	1	2%	80	3%	11	3%	45	2%	9	4%	-	-
Male-to-male sexual contact & IDU & Heterosexual contact	-	-	-	-	9	0%	-	-	23	1%	3	1%	-	-
Perinatal exposure	-	-	5	10%	31	1%	8	2%	7	0%	4	2%	-	-
Other/unknown	-	-	17	35%	437	18%	47	14%	324	12%	32	12%	21	95%
Total	6		49		2,427		335		2,775		257		22	

Notes:

Living with diagnosed HIV infection represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31, 2018. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31, 2018, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated because census data unavailable or for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

^b The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^c Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^d Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and selected characteristics, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2014-2018

Characteristic	Living with Diagnosed HIV Infection														
	2014			2015			2016			2017			2018		
	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%	Rate ^a	No.	%
Sex at birth															
Males	451.2	4,281	82%	451.7	4,347	81%	460.0	4,475	81%	466.0	4,617	81%	465.6	4,680	80%
Females	97.4	956	18%	102.4	1,019	19%	105.8	1,063	19%	108.9	1,114	19%	114.9	1,191	20%
Age at end of year															
<13	5.3	18	<1%	5.6	19	<1%	5.0	17	<1%	4.3	15	<1%	4.3	15	<1%
13-14	11.6	6	<1%	9.7	5	<1%	9.8	5	<1%	9.7	5	<1%	9.5	5	<1%
15-19	17.3	22	<1%	17.8	23	<1%	24.6	32	1%	23.3	31	1%	21.7	29	<1%
20-24	194.1	259	5%	164.7	216	4%	135.9	178	3%	128.0	170	3%	121.3	162	3%
25-29	305.8	475	9%	312.1	501	9%	308.9	507	9%	328.8	547	10%	317.0	531	9%
30-34	343.8	490	9%	344.6	499	9%	360.7	533	10%	370.7	562	10%	395.4	622	11%
35-39	400.9	523	10%	413.2	554	10%	435.5	600	11%	438.0	621	11%	419.2	609	10%
40-44	506.4	671	13%	472.5	617	11%	457.3	581	10%	452.0	576	10%	446.5	577	10%
45-49	690.2	883	17%	660.9	860	16%	617.4	818	15%	571.3	766	13%	565.5	755	13%
50-54	642.2	862	16%	675.7	901	17%	720.5	936	17%	734.9	941	16%	742.4	940	16%
55-64	370.8	846	16%	406.0	950	18%	446.7	1,060	19%	489.8	1,184	21%	516.5	1,265	22%
65+	79.5	182	3%	93.1	221	4%	110.3	271	5%	121.3	313	5%	134.6	361	6%
Race/Ethnicity^b															
American Indian/Alaska Native	128.5	5	<1%	125.9	5	<1%	126.1	5	<1%	151.1	6	<1%	150.7	6	<1%
Asian/Pacific Islander	40.1	29	1%	44.1	34	1%	50.4	42	1%	50.7	46	1%	49.7	49	1%
Black/African-American	670.9	1,992	38%	677.0	2,066	39%	699.6	2,191	40%	716.0	2,327	41%	727.0	2,427	41%
Hispanic/Latinx	355.3	271	5%	350.1	276	5%	357.9	292	5%	365.8	314	5%	369.1	335	6%
White	187.1	2,685	51%	188.0	2,717	51%	189.3	2,738	49%	189.7	2,765	48%	189.7	2,775	47%
Multi-Race	514.4	233	4%	522.7	246	5%	506.2	248	4%	494.6	251	4%	489.8	257	4%
Unknown	*	22	<1%	*	22	<1%	*	22	<1%	*	22	<1%	*	22	<1%
Race/Ethnicity^b and Sex at birth															
American Indian/Alaska Native Males	*	4	<1%	*	4	<1%	*	4	<1%	253.0	5	<1%	252.5	5	<1%
American Indian/Alaska Native Females	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%	*	1	<1%
Asian/Pacific Islander Males	67.8	24	<1%	68.8	26	<1%	77.9	32	1%	78.7	35	1%	74.4	36	1%
Asian/Pacific Islander Females	13.5	5	<1%	20.3	8	<1%	23.6	10	<1%	23.7	11	<1%	25.9	13	<1%
Black/African-American Males	975.0	1,393	27%	966.0	1,420	26%	999.3	1,512	27%	1,022.8	1,608	28%	1,024.2	1,659	28%
Black/African-American Females	388.8	599	11%	408.4	646	12%	419.5	679	12%	428.6	719	13%	446.9	768	13%
Hispanic/Latino Males	532.8	212	4%	528.2	216	4%	545.3	230	4%	564.6	251	4%	573.7	271	5%
Hispanic/Latina Females	161.7	59	1%	158.1	60	1%	157.4	62	1%	152.2	63	1%	147.0	64	1%
White Males	344.1	2,432	46%	345.1	2,457	46%	347.0	2,471	45%	347.2	2,492	43%	344.8	2,482	42%
White Females	34.7	253	5%	35.5	260	5%	36.4	267	5%	36.9	273	5%	39.5	293	5%
Multi-Race Males	898.3	198	4%	899.0	206	4%	864.6	208	4%	833.2	208	4%	809.5	209	4%
Multi-Race Females	150.5	35	1%	165.7	40	1%	160.4	40	1%	166.8	43	1%	180.1	48	1%
Unknown	*	22	<1%	*	22	<1%	*	22	<1%	*	22	<1%	*	22	<1%
Total	271.3	5,237		274.2	5,366		280.1	5,538		284.5	5,731		287.5	5,871	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count <5 due to unstable rates. Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with diagnosed HIV infection per 100,000 population calculated using U.S. Census estimates for that year.

^b Hispanics/Latinx may be of any race. Persons with a race of American Indian/Alaska Native, Asian/Pacific Islander, Black/African-American, White, or Multi-race are not-Hispanic. Asian/Pacific Islander includes Native Hawaiians.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and transmission category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2014-2018

Living with Diagnosed HIV Infection										
Transmission Category ^a	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent										
Male-to-male sexual contact	3,103	73%	3,136	72%	3,214	72%	3,299	72%	3,335	72%
Injection drug use (IDU)	96	2%	93	2%	100	2%	107	2%	111	2%
Male-to-male sexual contact and IDU	182	4%	186	4%	196	4%	206	4%	195	4%
Heterosexual contact	239	6%	236	5%	242	5%	241	5%	256	5%
Other/unknown	641	15%	675	16%	700	16%	740	16%	758	16%
Subtotal	4,261	100%	4,326	100%	4,452	100%	4,593	100%	4,655	100%
Female adult or adolescent										
Injection drug use	84	9%	93	9%	95	9%	99	9%	109	9%
Heterosexual contact	738	80%	788	80%	827	80%	874	81%	931	80%
Other/unknown	104	11%	107	11%	110	11%	110	10%	119	10%
Subtotal	926	100%	988	100%	1,032	100%	1,083	100%	1,159	100%
Child (<13 yrs at diagnosis)										
Perinatal	44	88%	46	88%	48	89%	50	91%	52	91%
Other/unknown	6	12%	6	12%	6	11%	5	9%	5	9%
Subtotal	50	100%	52	100%	54	100%	55	100%	57	100%
Total	5,237		5,366		5,538		5,731		5,871	

Trends in reported persons living with diagnosed HIV infection by year of diagnosis and exposure category, HIV Prevention Planning Region 11/Ryan White Part A-Columbus, 2014-2018

Living with Diagnosed HIV Infection										
Exposure Category ^b	2014		2015		2016		2017		2018	
	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact only	2,941	56%	2,979	56%	3,058	55%	3,140	55%	3,189	54%
Injection drug use (IDU) only	86	2%	89	2%	92	2%	95	2%	108	2%
Heterosexual contact only	978	19%	1,025	19%	1,070	19%	1,116	19%	1,188	20%
Male-to-male sexual contact & IDU	147	3%	149	3%	157	3%	167	3%	160	3%
IDU & Heterosexual contact	94	2%	97	2%	103	2%	111	2%	112	2%
Male-to-male sexual contact & Heterosexual contact	162	3%	157	3%	156	3%	159	3%	146	2%
Male-to-male sexual contact & IDU & Heterosexual contact	35	1%	37	1%	39	1%	39	1%	35	1%
Perinatal exposure	46	1%	48	1%	51	1%	53	1%	55	1%
Other/unknown	748	14%	785	15%	812	15%	851	15%	878	15%
Total	5,237		5,366		5,538		5,731		5,871	

Notes:

Living with diagnosed HIV infection by year (2014-2018) represents all persons ever diagnosed and reported with HIV and/or AIDS who have not been reported as having died as of December 31 of the corresponding year. Persons living with diagnosed HIV infection represent persons living in Ohio as of December 31 of the corresponding year, regardless of whether the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category.

^a Transmission categories are mutually exclusive, hierarchical risk categories determined by the CDC and system-calculated using sex at birth and risk factor history to determine mode of transmission. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm. Thus, transgender women are included in the male-to-male sexual contact transmission category if assigned male at birth and risk factor history indicates sex with males. Please note this is for the categorization of HIV transmission categories only and not to describe sexual orientation.

^b Exposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health, HIV/AIDS Surveillance Program. Data reported through June 30, 2019.